

## Author Index

- Åkesson, K., Nyström, L., Färnkvist, L., Östman, J., Lernmark, Å. & Kockum, I. and the DISS study group Increased risk of diabetes among relatives of female insulin-treated patients diagnosed at 15–34 years of age, 1551
- Åman, J. *see* Särnblad, S.
- Aas, A.M., Bergstad, I., Thorsby, P.M., Johannesen, Ø., Solberg, M. & Birkeland, K.I. An intensified lifestyle intervention programme may be superior to insulin treatment in poorly controlled Type 2 diabetic patients on oral hypoglycaemic agents: results of a feasibility study, 316
- Abbas, M. *see* Abdelgadir, M.
- Abbas, Z.G., Lutale, J. & Archibald, L.K. Rodent bites on the feet of diabetes patients in Tanzania, 631
- Abdelgadir, M., Abbas, M., Järvi, A., Elbagir, M., Eltom, M. & Berne, C. Glycaemic and insulin responses of six traditional Sudanese carbohydrate-rich meals in subjects with Type 2 diabetes mellitus, 213
- Abdella, N.A., Mojiminiyi, O.A., Moussa, M.A., Zaki, M., Al Mohammedi, H., Al Ozairi, E.S.S. & Al Jebely, S. Plasma leptin concentration in patients with Type 2 diabetes: relationship to cardiovascular disease risk factors and insulin resistance, 278
- Abletshauser, C. *see* Bengel, F.M.
- Acosta, J.A. *see* Rutkove, S.B.
- Adaikalakoteswari, A., Balasubramanyam, M. & Mohan, V. Telomere shortening occurs in Asian Indian Type 2 diabetic patients, 1151
- Adams, J., Pearce, M.S., White, M., Unwin, N.C. & Parker, L. No consistent association between birthweight and parental risk of diabetes and cardiovascular disease, 950
- Adams, L.A. & Angulo, P. Recent concepts in non-alcoholic fatty liver disease, 1129
- Adler, A.I., Levy, J.C., Matthews, D.R., Stratton, I.M., Hines, G. & Holman, R.R. Insulin sensitivity at diagnosis of Type 2 diabetes is not associated with subsequent cardiovascular disease (UKPDS 67), 306
- ADVANCE Collaborative Group. ADVANCE—Action in Diabetes and Vascular Disease: patient recruitment and characteristics of the study population at baseline, 882
- Advani, A. & Taylor, R. Life-threatening hypokalaemia on a low-carbohydrate diet associated with previously undiagnosed primary hypoadosteronism, 1605
- Advani, A. Handbook of Diabetes, 3rd edn, 817
- Agardh, C.-D. *see* Torffvit, O.
- Agardh, E. *see* Torffvit, O.
- Agarwal, M.M., Dhath, G.S., Punnose, J. & Koster, G. Gestational diabetes: dilemma caused by multiple international diagnostic criteria, 1731
- Agewall, S. *see* Henareh, L.
- Aguilera, E., Casamitjana, R., Ercilla, G., Oriola, J., Nicoletti, F., Gomis, R. & Conget, I. Clinical characteristics,  $\beta$ -cell function, HLA class II and mutations in MODY genes in non-paediatric subjects with Type 1 diabetes without pancreatic autoantibodies, 137
- Ahmed, A.T. *see* Karter, A.J.
- Ahn, Y.-B. *see* Ko, S.-H.
- Ainla, T., Baburin, A., Teesalu, R. & Rahu, M. The association between hyperglycaemia on admission and 180-day mortality in acute myocardial infarction patients with and without diabetes, 1321
- Airey, F.L. *see* McMahon, S.K.
- Ajayi, A.A.L. Gender influences the impact of sickle cell trait on Type 2 diabetic complications, 954
- Al Jebely, S. *see* Abdella, N.A.
- Al Mohammedi, H. *see* Abdella, N.A.
- Al Ozairi, E.S.S. *see* Abdella, N.A.
- Albayram, S. *see* Ozer, H.
- Al-Bermani, A., Desha, Y.H., Morgan, J., Soobrah, R., Symonds, C.S. & Taylor, R. Management of incidental hyperglycaemia in acute medical emergencies, 937
- Alberti, K.G.M.M. *see* Söderberg, S.
- Alberti, K.G.M.M. *see* Whincup, P.H.
- Alcolado, J.C. *see* Rees, D.A.
- Aldington, S.J. *see* Schneider, S.
- Ali, S.M.K. *see* Hussain, A.
- Allard, S.E. *see* Wotherspoon, F.
- Allen, S.E. *see* Douek, I.F.
- Allen-Taylor, L. *see* Margolis, D.J.
- Alviggi, L. *see* Seghieri, G.
- Ambye, L. *see* Bing, C.
- Amiel, S.A. *see* Rodrigues, I.A.S.
- An, B. *see* Feinglos, M.N.
- Andersen, G. *see* Rose, C.S.
- Andersen, H., Schmitz, O. & Nielsen, S. Decreased isometric muscle strength after acute hyperglycaemia in Type 1 diabetic patients, 1401
- Andersen, N.H., Poulsen, S.H., Knudsen, S.T., Heickendorff, L. & Mogensen, C.E. NT-proBNP in normoalbuminuric patients with Type 2 diabetes mellitus, 188
- Andersen, N.H., Poulsen, S.H., Poulsen, P.L., Knudsen, S.T., Helleberg, K., Hansen, K.W., Berg, T.J., Flyvbjerg, A. & Mogensen, C.E. Left ventricular dysfunction in hypertensive patients with Type 2 diabetes mellitus, 1218
- Anderson, P.J. *see* Northam, E.A.
- Andersson, S.A. *see* Lawlor, D.A.
- Andersson, S. *see* Loukovaara, M.
- Andreasen, A.H. *see* Nordly, S.
- Angulo, P. *see* Adams, L.A.
- Anichini, R. *see* Seghieri, G.
- Apostolou, O. *see* Sotiropoulos, A.
- Archibald, L.K. *see* Abbas, Z.G.
- Ardawi, M.S.M. *see* Damanhour, L.H.

## Author Index

- Åkesson, K., Nyström, L., Färnkvist, L., Östman, J., Lernmark, Å. & Kockum, I. and the DISS study group Increased risk of diabetes among relatives of female insulin-treated patients diagnosed at 15–34 years of age, 1551
- Åman, J. *see* Särnblad, S.
- Aas, A.M., Bergstad, I., Thorsby, P.M., Johannesen, Ø., Solberg, M. & Birkeland, K.I. An intensified lifestyle intervention programme may be superior to insulin treatment in poorly controlled Type 2 diabetic patients on oral hypoglycaemic agents: results of a feasibility study, 316
- Abbas, M. *see* Abdelgadir, M.
- Abbas, Z.G., Lutale, J. & Archibald, L.K. Rodent bites on the feet of diabetes patients in Tanzania, 631
- Abdelgadir, M., Abbas, M., Järvi, A., Elbagir, M., Eltom, M. & Berne, C. Glycaemic and insulin responses of six traditional Sudanese carbohydrate-rich meals in subjects with Type 2 diabetes mellitus, 213
- Abdella, N.A., Mojiminiyi, O.A., Moussa, M.A., Zaki, M., Al Mohammedi, H., Al Ozairi, E.S.S. & Al Jebely, S. Plasma leptin concentration in patients with Type 2 diabetes: relationship to cardiovascular disease risk factors and insulin resistance, 278
- Abletshauser, C. *see* Bengel, F.M.
- Acosta, J.A. *see* Rutkove, S.B.
- Adaikalakoteswari, A., Balasubramanyam, M. & Mohan, V. Telomere shortening occurs in Asian Indian Type 2 diabetic patients, 1151
- Adams, J., Pearce, M.S., White, M., Unwin, N.C. & Parker, L. No consistent association between birthweight and parental risk of diabetes and cardiovascular disease, 950
- Adams, L.A. & Angulo, P. Recent concepts in non-alcoholic fatty liver disease, 1129
- Adler, A.I., Levy, J.C., Matthews, D.R., Stratton, I.M., Hines, G. & Holman, R.R. Insulin sensitivity at diagnosis of Type 2 diabetes is not associated with subsequent cardiovascular disease (UKPDS 67), 306
- ADVANCE Collaborative Group. ADVANCE—Action in Diabetes and Vascular Disease: patient recruitment and characteristics of the study population at baseline, 882
- Advani, A. & Taylor, R. Life-threatening hypokalaemia on a low-carbohydrate diet associated with previously undiagnosed primary hypoadosteronism, 1605
- Advani, A. Handbook of Diabetes, 3rd edn, 817
- Agardh, C.-D. *see* Torffvit, O.
- Agardh, E. *see* Torffvit, O.
- Agarwal, M.M., Dhath, G.S., Punnose, J. & Koster, G. Gestational diabetes: dilemma caused by multiple international diagnostic criteria, 1731
- Agewall, S. *see* Henareh, L.
- Aguilera, E., Casamitjana, R., Ercilla, G., Oriola, J., Nicoletti, F., Gomis, R. & Conget, I. Clinical characteristics,  $\beta$ -cell function, HLA class II and mutations in MODY genes in non-paediatric subjects with Type 1 diabetes without pancreatic autoantibodies, 137
- Ahmed, A.T. *see* Karter, A.J.
- Ahn, Y.-B. *see* Ko, S.-H.
- Ainla, T., Baburin, A., Teesalu, R. & Rahu, M. The association between hyperglycaemia on admission and 180-day mortality in acute myocardial infarction patients with and without diabetes, 1321
- Airey, F.L. *see* McMahon, S.K.
- Ajayi, A.A.L. Gender influences the impact of sickle cell trait on Type 2 diabetic complications, 954
- Al Jebely, S. *see* Abdella, N.A.
- Al Mohammedi, H. *see* Abdella, N.A.
- Al Ozairi, E.S.S. *see* Abdella, N.A.
- Albayram, S. *see* Ozer, H.
- Al-Bermani, A., Desha, Y.H., Morgan, J., Soobrah, R., Symonds, C.S. & Taylor, R. Management of incidental hyperglycaemia in acute medical emergencies, 937
- Alberti, K.G.M.M. *see* Söderberg, S.
- Alberti, K.G.M.M. *see* Whincup, P.H.
- Alcolado, J.C. *see* Rees, D.A.
- Aldington, S.J. *see* Schneider, S.
- Ali, S.M.K. *see* Hussain, A.
- Allard, S.E. *see* Wotherspoon, F.
- Allen, S.E. *see* Douek, I.F.
- Allen-Taylor, L. *see* Margolis, D.J.
- Alviggi, L. *see* Seghieri, G.
- Ambye, L. *see* Bing, C.
- Amiel, S.A. *see* Rodrigues, I.A.S.
- An, B. *see* Feinglos, M.N.
- Andersen, G. *see* Rose, C.S.
- Andersen, H., Schmitz, O. & Nielsen, S. Decreased isometric muscle strength after acute hyperglycaemia in Type 1 diabetic patients, 1401
- Andersen, N.H., Poulsen, S.H., Knudsen, S.T., Heickendorff, L. & Mogensen, C.E. NT-proBNP in normoalbuminuric patients with Type 2 diabetes mellitus, 188
- Andersen, N.H., Poulsen, S.H., Poulsen, P.L., Knudsen, S.T., Helleberg, K., Hansen, K.W., Berg, T.J., Flyvbjerg, A. & Mogensen, C.E. Left ventricular dysfunction in hypertensive patients with Type 2 diabetes mellitus, 1218
- Anderson, P.J. *see* Northam, E.A.
- Andersson, S.A. *see* Lawlor, D.A.
- Andersson, S. *see* Loukovaara, M.
- Andreasen, A.H. *see* Nordly, S.
- Angulo, P. *see* Adams, L.A.
- Anichini, R. *see* Seghieri, G.
- Apostolou, O. *see* Sotiropoulos, A.
- Archibald, L.K. *see* Abbas, Z.G.
- Ardawi, M.S.M. *see* Damanhour, L.H.

- Ardill, J. *see* Lindsay, J.R.  
 Arii, K. *see* Ikeda, Y.  
 Armour, T.A., Norris, S.L., Jack Jr, L. Zhang, X. & Fisher, L. The effectiveness of family interventions in people with diabetes mellitus: a systematic review, 1295  
 Armstrong, N. *see* Middlebrooke, A.R.  
 Aronson, D. & Levy, Y. Reply, 509  
 Arstila, M. *see* Pajunen, P.  
 Arteagoitia, J.M. *see* Larrañaga, I.  
 Assal, J.-P. *see* Pataky, Z.  
 Astrup, A. *see* Black, E.  
 Atkin, S.L. *see* González, S.  
 Atkinson, A.B. *see* Au, S.  
 Atkinson, A.B. *see* McSorley, P.T.  
 Atkinson, M. *see* Gale, E.A.M.  
 Attali, J.R. *see* N'Guyen Thi, N.  
 Au, S., Courtney, C.H., Ennis, C.N., Sheridan, B., Atkinson, A.B. & Bell, P.M. The effect of manipulation of basal pulsatile insulin on insulin action in Type 2 diabetes, 1064  
 Augendre-Ferrante, B. *see* Malone, J.K.  
 Avogaro, A. *see* Perriello, G.  
 Aydin, A. *see* Szelestei, J.  
 Azad Khan, A.K. *see* Hussain, A.  
 Azad Khan, A.K. *see* Sayeed, M.A.
- Babu, H.N.S. *see* Vimalaswaran, K.S.  
 Baburin, A. *see* Ainla, T.  
 Bader, M.S., Hinthorn, D., Lai, S.M. & Ellerbeck, E.F. Hyperglycaemia and mortality of diabetic patients with candidaemia, 1252  
 Bähring, S. *see* Szelestei, J.  
 Bai, S. *see* Malone, J.K.  
 Baillet-Blanco, L. *see* Rigalleau, V.  
 Bain, L. *see* Royle, P.L.  
 Bakiner, O. *see* Shain, M.  
 Bakri, D. *see* Shehadeh, N.  
 Balasubramanyam, A. *see* Maldonado, M.R.  
 Balasubramanyam, M. *see* Adaikalakoteswari, A.  
 Ball, V. *see* Roland, J.M.  
 Band, M.M. *see* Donnelly, L.A.  
 Banerjee, S. *see* Denniston, A.K.O.  
 Banu, A. *see* Sayeed, M.A.  
 Bardini, G. *see* Mannucci, E.  
 Barkai, L., Soós, A. & Vámosi, I. Association of angiotensin-converting enzyme DD genotype with 24-h blood pressure abnormalities in normoalbuminuric children and adolescents with Type 1 diabetes, 1054  
 Barmada, M.M. *see* Schneider, A.  
 Barnett, A.H. *see* Patel, A.  
 Barrett, T.G. *see* Kershaw, M.J.R.  
 Barrie, P. *see* Sampson, M.J.  
 Barroso, I. Genetics of Type 2 diabetes, 517  
 Barsotti, M. *see* Giannarelli, R.  
 Barthel, A. *see* Schinner, S.  
 Barton, D.M. *see* Buch, H.N.  
 Battler, A. *see* Dotevall, A.  
 Baumann, M.E. *see* Sagen, J.V.  
 Bayegan, K. *see* Frossard, M.  
 Bebb, C., Kendrick, D., Stewart, J., Coupland, C., Madeley, R., Brown, K., Burden, R. & Sturrock, N. for the Nottingham Diabetes Blood Pressure Study Inequalities in glycaemic control in patients with Type 2 diabetes in primary care, 1364  
 Beck, M.J. *see* O'Bryan, T.A.  
 Becker, D. *see* Bosnyak, Z.  
 Beck-Nielsen, H. *see* Hoffmann, S.K.S.  
 Beeching, N.J. *see* Evans, E.M.  
 Beekman, A.T. *see* Pouwer, F.  
 Begic, D. *see* Pibernik-Okanovic, M.  
 Begum, R. *see* Sayeed, M.A.  
 Belch, J.J.F. *see* Newton, D.J.  
 Belcher, G. & Scherthaner, G. Changes in liver tests during 1-year treatment of patients with Type 2 diabetes with pioglitazone, metformin or gliclazide, 973  
 Belfiglio, M. *see* Franciosi, M.  
 Belicza, M. *see* Mlinaric Missoni, E.  
 Bell, P.M. *see* Au, S.  
 Bell, P.M. *see* Lindsay, J.R.  
 Bell, P.M. *see* McSorley, P.T.  
 Benbow, S.J. *see* Daousi, C.  
 Bengel, F.M., Abletshauser, C., Neverve, J., Schnell, O., Nekolla, S.G., Standl, E. & Schwaiger, M. Effects of nateglinide on myocardial microvascular reactivity in Type 2 diabetes mellitus – a randomized study using positron emission tomography, 158  
 Bennett, C.E. *see* Carter, A.M.  
 Bennett, S.M.A., Heise, M., Jones, N.P., Walker, M. & Wilding, J.P.H. Reply, 667  
 Bennett, S.P. *see* Newton, D.J.  
 Bennetts, B. *see* Donaghue, K.C.  
 Bentzen, J. *see* Koster, B.  
 Benvenuto, A. *see* Caselli, A.  
 Benzie, I.F.F. *see* Choi, S.W.  
 Berg, T.J. *see* Andersen, N.H.  
 Bergstad, I. *see* Aas, A.M.  
 Bergstrom, C. *see* Linfoot, P.  
 Berla, M. *see* Rachmani, R.  
 Berlin, J.A. *see* Margolis, D.J.  
 Bermejo, E. *see* Martínez-Frías, M.L.  
 Berne, C. *see* Abdelgadir, M.  
 Berne, C. on behalf of the Orlistat Swedish Type 2 diabetes Study Group. A randomized study of orlistat in combination with a weight management programme in obese patients with Type 2 diabetes treated with metformin, 612  
 Berry, K. *see* Kershaw, M.J.R.  
 Bertolini, L. *see* Targher, G.  
 Best, J.D. *see* Schutte, A.E.  
 Bets, D. *see* Wulffélé, M.G.  
 Betts, P., Mulligan, J., Ward, P., Smith, B. & Wilkin, T. Increasing body weight predicts the earlier onset of insulin-dependant diabetes in childhood: testing the 'accelerator hypothesis' (2), 144  
 Biddulph, J.P. *see* Thomason, M.J.  
 Bing, C., Ambye, L., Fenger, M., Jorgensen, T., Borch-Johnsen, K., Madbsad, S. & Urhammer, S.A. Large-scale studies of the Leu72Met polymorphism of the ghrelin gene in relation to the metabolic syndrome and associated quantitative traits, 1157  
 Bingley, P.J. for the Metformin Trial Group *see* Douek, I.F.  
 Birkeland, K.I. *see* Aas, A.M.  
 Bisseling, T.M., Wouterse, A.C., Steegers, E.A.P., Elving, L., Russel, F.G.M. & Smits, P. Dysfunction of the cyclo-oxygenase pathway in the foetoplacental circulation in Type 1 diabetes mellitus, 503

Ryan C	Solomon D	Tong PC	Westaway M
Ryan E	Songer T	Toplak H	Weston P
Ryder E	Songini M	Tossavainen P	Wexler D
Sajadieh A	Spangler J	Trauner M	Whincup P
Saletsky R	Speight J	Tringham J	White K
Sami S	Spijkerman A	Troughton J	White N
Sanchez-Albisua I	Spranger J	Tsatsoulis A	Whitford D
Sandbaek A	Squirrell D	Tuomilehto J	Widhalm K
Sandhu M	Stefanidis A	Tura A	Wiener K
Sarnblad S	Steffes MW	Turnheim K	Wierzbicki A
Sattar N	Stehouwer C	Turunen J	Wild S
Saunders S	Stephens J	Twigg S	Wilder R
Savage D	Stern M	Umpierrez G	Wilding J
Savage M	Stevens MJ	Vaccaro O	Wilkinson IB
Sayeed M	Stevens P	Vaidya A	Williams A
Scarpello JHB	Stevens R	Vaidya B	Williams B
Schaper N	Stewart J	Valabhji J	Winter W
Schatz D	Stingl H	Valensi P	Witte D
Schatz H	Stirban A	Valentine W	Wittert G
Schectman J	Stolk RP	Van Belle E	Wolever T
Scherthaner G	Strachan MWJ	van Bellen B	Wolfsdorf J
Schlott N	Stratton I	Van Haefen T	Wollesen F
Schlossman D	Stuart L	Vanschic C	Wong K
Schmitz O	Stumvoll M	Vardi Y	Wong S
Schooling M	Sturrock NDC	Vasseur F	Woo J
Schulze M	Sturt J	vaughan	Woo KS
Scott A	Sudhop T	Vella A	Woo V
Scuffham P	Suminski R	Vendrell J	Woodcock A
Scuteri A	Sundkvist G	Vinik A	Woods D
Sedor J	Svensson J	Vionnet N	Wu A
Seghieri G	Svensson M	Viswanthan N	Wu S
Selby PL	Swift PGF	Vitale C	Yale J-F
Selvin E	Tack C	Vohl M-C	Yamamoto S
Senior P	Tait K	Vora J	Yang B
Sesti G	Takebayashi K	Wackers F	Yang X
Seukeran D	Tamborlane W	Wagner A	Yee M
Shaban C	Tapp R	Wahid S	Yildiz A
Shaw J	Targher G	Wahlberg E	Yilmaz M
Shiwaku K	Tarnow L	Wales JKH	Yin W
Shore A	Tattersall RB	Walker J	Yngen M
shotliff	Taylor R	Walker M	Yokoyama H
Sibal L	Taylor R	Walker M	You T
Siebenhofer A	Tedesco MA	Walkinshaw S	Young IS
Siegrist J	Tekeli A	Walton C	Young M
Siemianowicz K	Temple RC	Wannamethee G	Young R
Silveiro S	Tesfaye S	Warren R	Younis N
Simmons D	Thomas A	Wascher T	Yu TS
Simsek S	Thomas R	Watts G	Yumuk Z
Sinclair A	Thomas S	Weaver J	Ziegler A-G
Singh B	Thomas T	Webber J	Zimny S
Sjoholm A	Thornalley P	Weedon M	Zingg J-M
Sjolie AK	Tikellis G	Weinger K	Zinken K
Skinner C	Tillin T	Welters H	Zoffmann V
Slart RH	Tindall H	Wens J	
Sochett E	Tomiyama H	Wessels A	



- Ardill, J. *see* Lindsay, J.R.  
 Arie, K. *see* Ikeda, Y.  
 Armour, T.A., Norris, S.L., Jack Jr, L. Zhang, X. & Fisher, L. The effectiveness of family interventions in people with diabetes mellitus: a systematic review, 1295  
 Armstrong, N. *see* Middlebrooke, A.R.  
 Aronson, D. & Levy, Y. Reply, 509  
 Arstila, M. *see* Pajunen, P.  
 Arteagoitia, J.M. *see* Larrañaga, I.  
 Assal, J.-P. *see* Pataky, Z.  
 Astrup, A. *see* Black, E.  
 Atkin, S.L. *see* González, S.  
 Atkinson, A.B. *see* Au, S.  
 Atkinson, A.B. *see* McSorley, P.T.  
 Atkinson, M. *see* Gale, E.A.M.  
 Attali, J.R. *see* N'Guyen Thi, N.  
 Au, S., Courtney, C.H., Ennis, C.N., Sheridan, B., Atkinson, A.B. & Bell, P.M. The effect of manipulation of basal pulsatile insulin on insulin action in Type 2 diabetes, 1064  
 Augendre-Ferrante, B. *see* Malone, J.K.  
 Avogaro, A. *see* Perriello, G.  
 Aydin, A. *see* Szelestei, J.  
 Azad Khan, A.K. *see* Hussain, A.  
 Azad Khan, A.K. *see* Sayeed, M.A.  
 Babu, H.N.S. *see* Vimalaswaran, K.S.  
 Baburin, A. *see* Ainla, T.  
 Bader, M.S., Hinthorn, D., Lai, S.M. & Ellerbeck, E.F. Hyperglycaemia and mortality of diabetic patients with candidaemia, 1252  
 Bähring, S. *see* Szelestei, J.  
 Bai, S. *see* Malone, J.K.  
 Baillet-Blanco, L. *see* Rigalleau, V.  
 Bain, L. *see* Royle, P.L.  
 Bakiner, O. *see* Shain, M.  
 Bakri, D. *see* Shehadeh, N.  
 Balasubramanyam, A. *see* Maldonado, M.R.  
 Balasubramanyam, M. *see* Adaikalakoteswari, A.  
 Ball, V. *see* Roland, J.M.  
 Band, M.M. *see* Donnelly, L.A.  
 Banerjee, S. *see* Denniston, A.K.O.  
 Banu, A. *see* Sayeed, M.A.  
 Bardini, G. *see* Mannucci, E.  
 Barkai, L., Soós, A. & Vámosi, I. Association of angiotensin-converting enzyme DD genotype with 24-h blood pressure abnormalities in normoalbuminuric children and adolescents with Type 1 diabetes, 1054  
 Barmada, M.M. *see* Schneider, A.  
 Barnett, A.H. *see* Patel, A.  
 Barrett, T.G. *see* Kershaw, M.J.R.  
 Barrie, P. *see* Sampson, M.J.  
 Barroso, I. Genetics of Type 2 diabetes, 517  
 Barsotti, M. *see* Giannarelli, R.  
 Barthel, A. *see* Schinner, S.  
 Barton, D.M. *see* Buch, H.N.  
 Battler, A. *see* Dotevall, A.  
 Baumann, M.E. *see* Sagen, J.V.  
 Bayegan, K. *see* Frossard, M.  
 Bebb, C., Kendrick, D., Stewart, J., Coupland, C., Madeley, R., Brown, K., Burden, R. & Sturrock, N. for the Nottingham Diabetes Blood Pressure Study Inequalities in glycaemic control in patients with Type 2 diabetes in primary care, 1364  
 Beck, M.J. *see* O'Bryan, T.A.  
 Becker, D. *see* Bosnyak, Z.  
 Beck-Nielsen, H. *see* Hoffmann, S.K.S.  
 Beeching, N.J. *see* Evans, E.M.  
 Beekman, A.T. *see* Pouwer, F.  
 Begic, D. *see* Pibernik-Okanovic, M.  
 Begum, R. *see* Sayeed, M.A.  
 Belch, J.J.F. *see* Newton, D.J.  
 Belcher, G. & Scherthaner, G. Changes in liver tests during 1-year treatment of patients with Type 2 diabetes with pioglitazone, metformin or gliclazide, 973  
 Belfiglio, M. *see* Franciosi, M.  
 Beliczka, M. *see* Mlinaric Missoni, E.  
 Bell, P.M. *see* Au, S.  
 Bell, P.M. *see* Lindsay, J.R.  
 Bell, P.M. *see* McSorley, P.T.  
 Benbow, S.J. *see* Daousi, C.  
 Bengel, F.M., Abletshauser, C., Neverve, J., Schnell, O., Nekolla, S.G., Standl, E. & Schwaiger, M. Effects of nateglinide on myocardial microvascular reactivity in Type 2 diabetes mellitus – a randomized study using positron emission tomography, 158  
 Bennett, C.E. *see* Carter, A.M.  
 Bennett, S.M.A., Heise, M., Jones, N.P., Walker, M. & Wilding, J.P.H. Reply, 667  
 Bennett, S.P. *see* Newton, D.J.  
 Bennets, B. *see* Donaghue, K.C.  
 Bentzen, J. *see* Koster, B.  
 Benvenuto, A. *see* Caselli, A.  
 Benzie, I.F.F. *see* Choi, S.W.  
 Berg, T.J. *see* Andersen, N.H.  
 Bergstad, I. *see* Aas, A.M.  
 Bergstrom, C. *see* Linfoot, P.  
 Berla, M. *see* Rachmani, R.  
 Berlin, J.A. *see* Margolis, D.J.  
 Bermejo, E. *see* Martínez-Frías, M.L.  
 Berne, C. *see* Abdelgadir, M.  
 Berne, C. on behalf of the Orlistat Swedish Type 2 diabetes Study Group. A randomized study of orlistat in combination with a weight management programme in obese patients with Type 2 diabetes treated with metformin, 612  
 Berry, K. *see* Kershaw, M.J.R.  
 Bertolini, L. *see* Targher, G.  
 Best, J.D. *see* Schutte, A.E.  
 Bets, D. *see* Wulffélé, M.G.  
 Betts, P., Mulligan, J., Ward, P., Smith, B. & Wilkin, T. Increasing body weight predicts the earlier onset of insulin-dependant diabetes in childhood: testing the 'accelerator hypothesis' (2), 144  
 Biddulph, J.P. *see* Thomason, M.J.  
 Bing, C., Ambye, L., Fenger, M., Jørgensen, T., Borch-Johnsen, K., Madshad, S. & Urhammer, S.A. Large-scale studies of the Leu72Met polymorphism of the ghrelin gene in relation to the metabolic syndrome and associated quantitative traits, 1157  
 Bingley, P.J. for the Metformin Trial Group *see* Douek, I.F.  
 Birkeland, K.I. *see* Aas, A.M.  
 Bisseling, T.M., Wouterse, A.C., Steegers, E.A.P., Elving, L., Russel, F.G.M. & Smits, P. Dysfunction of the cyclo-oxygenase pathway in the foetoplacental circulation in Type 1 diabetes mellitus, 503

- Bitó, T., Földesi, I., Nyári, T. & Pál, A. Prediction of gestational diabetes mellitus in a high-risk group by insulin measurement in early pregnancy, 1434
- Bitsanis, D. *see* Min, Y.
- Björkander, I. *see* Held, C.
- Black, E., Holst, C., Astrup, A., Toubro, S., Echwald, S., Pedersen, O. & Sørensen, T.I.A. Long-term influences of body-weight changes, independent of the attained weight, on risk of impaired glucose tolerance and Type 2 diabetes, 1199
- Blank, D. *see* Frossard, M.
- Blann, A.D. *see* Lim, H.S.
- Blaum, C.S., Volpato, S., Cappola, A.R., Chaves, P., Xue, Q.L., Guralnik, J.M. & Fried, L.P. Diabetes, hyperglycaemia and mortality in disabled older women: The Women's Health and Ageing Study I, 543
- Blom, H. *see* Veldman, B.A.J.
- Blomqvist, M. *see* Buschard, K.
- Bo, S., Ciccone, G., Rosato, R., Gancia, R., Grassi, G., Merletti, F. & Pagano, G.F. Renal damage in patients with Type 2 diabetes: a strong predictor of mortality, 258
- Bo, S., Gambino, R., Guidi, S., Silli, B., Gentile, L., Cassader, M. & Pagano, G.F. Plasma nitrotyrosine levels, antioxidant vitamins and hyperglycaemia, 1185
- Boaz, M. *see* Wainstein, J.
- Bochenski, C. *see* Wilson, C.
- Bodtger, U. & Wittrup, M. A rational clinical approach to suspected insulin allergy: status after five years and 22 cases, 102
- Boehm, B.O. *see* Karges, B.
- Bøg-Hansen, E. *see* Buschard, K.
- Boggi, U. *see* Giannarelli, R.
- Boland, L. *see* Selvin, E.
- Boldrini, A. *see* Di Cianni, G.
- Bonifacio, E. *see* Sordi, V.
- Bonomo, M., Corica, D., Mion, E., Gonçalves, D., Motta, G., Merati, R., Ragusa, A. & Morabito, A. Evaluating the therapeutic approach in pregnancies complicated by borderline glucose intolerance: a randomized clinical trial, 1536
- Borch-Johnsen, K. *see* Bing, C.
- Borch-Johnsen, K. *see* Glümer, C.
- Borch-Johnsen, K. *see* Hamid, Y.H.
- Borch-Johnsen, K. *see* Rose, C.S.
- Borch-Johnsen, K. *see* Sandbæk, A.
- Born, J. *see* Merl, V.
- Bornstein, S.R. *see* Schinner, S.
- Boronat, M., Chirino, R., Varillas, V.F., Saavedra, P., Marrero, D., Fábregas, M. & J. Nóvoa, F. Prevalence of the metabolic syndrome in the island of Gran Canaria: comparison of three major diagnostic proposals, 1751
- Bosi, E. *see* Perriello, G.
- Bosnyak, Z., Nishimura, R., Hagan Hughes, M., Tajima, N., Becker, D., Tuomilehto, J. & Orchard, T.J. Excess mortality in Black compared with White patients with Type 1 diabetes: an examination of underlying causes, 1636
- Bostock, J.A. *see* Carter, A.M.
- Bots, M.L. *see* Muis, M.J.
- Boucher, B.J., Timms, P.M. & Hitman, G.A. Association between fasting glucose and C-reactive protein in middle-aged subjects, 508
- Boyages, S.C. *see* Orr, N.J.
- Bradbury, S. *see* Buch, H.N.
- Brassøe, R., Elkmann, T., Hempel, M. & Gravholt, C.H. Fulminant lactic acidosis in two patients with Type 2 diabetes treated with metformin, 1451
- Bratthall, D. *see* Twetman, S.
- Brennan, A.M., McNamara, B., Plant, W.D. & O'Halloran, D.J. An atypical case of acute ischaemic monomelic neuropathy post vascular access surgery in a patient with Type 1 diabetes mellitus, 813
- Breschi, M.C. *see* Seghieri, G.
- Broch, M. *see* Vendrell, J.
- Brown, F. *see* Wyatt, J.W.
- Brown, K. *see* Bebb, C.
- Brown, R. *see* Glaab, L.A.
- Browne, D.L. *see* Wotherspoon, F.
- Bruce, D.G. *see* Davis, T.M.E.
- de Bruin, T.W.A. *see* van Dam, R.M.
- Brunetti, P. on behalf of the QUARTET Study Group *see* Charbonnel, B.H.
- Bruno, C.M., Cilio, D., Sciacca, C. & Pellicano, R. Circulating erythropoietin in diabetes mellitus, 1780
- Bu, H. *see* Ran, X.
- Buch, H.N., Barton, D.M., Varughese, G.I., Bradbury, S., Scarpello, J.H.B. & Walker, A.B. An assessment of the coverage of a district-wide diabetic retinopathy screening service, 840
- Buchanan, T.A. *see* Karim, R.
- Buonocore, M.H. *see* Glaser, N.S.
- Burden, R. *see* Bebb, C.
- Burke, J. *see* Coll, A.P.
- Burns, J.A. *see* McElduff, P.
- Buschard, K., Fredman, P., Bøg-Hansen, E., Blomqvist, M., Hedner, J., Råstam, L. & Lindblad, U. Low serum concentration of sulfatide and presence of sulfated lactosylceramid are associated with Type 2 diabetes. The Skaraborg Project, 1190
- But, H. *see* Ran, X.
- Buysschaert, M., Tomasi, J.P. & Hermans, M.P. Prospective screening for biopsy proven coeliac disease, autoimmunity and malabsorption markers in Belgian subjects with Type 1 diabetes, 889
- Byrne, C.D. *see* Guzder, R.N.
- Byrne, C.D. The Metabolic Syndrome: Diabetes, Obesity Hyperlipidemia and Hypertension, 513
- Cainarca, S. *see* Sordi, V.
- Caixàs, A. *see* Giménez-Pérez, G.
- Cameron, F.J. *see* Northam, E.A.
- Cameron, F.J., Kean, M.J., Wellard, R.M., Werther, G.A., Neil, J.J. & Inder, T.E. Insights into the acute cerebral metabolic changes associated with childhood diabetes, 648
- Campaigne, B.N. *see* Malone, J.K.
- Campbell, D.A. *see* Wraight, P.R.
- Canani, L.H. *see* Crispim, D.
- Canani, L.H., Costa, L.A., Crispim, D., Gonçalves dos Santos, K., Roisenberg, I., Lisbôa, H.R.K., Sarturi Tres, G., Maia, A.L. & Gross, J.L. The presence of allele D of angiotensin-converting enzyme polymorphism is associated with diabetic nephropathy in patients with less than 10 years duration of Type 2 diabetes, 1167
- Capaldo, B. *see* Di Bonito, P.
- Cappola, A.R. *see* Blaum, C.S.
- Caravaggi, C. *see* Faglia, E.
- Cardiovascular therapies and their role in diabetic eye disease, 665
- Cardona, F. *see* Morcillo, S.

- Cardwell, C.R., Carson, D.J. & Patterson, C.C. Parental age at delivery, birth order, birth weight and gestational age are associated with the risk of childhood Type 1 diabetes: a UK regional retrospective cohort study, 200
- Carlessi, R.M. *see* Crispim, D.
- Carne, C.L. *see* McMahon, S.K.
- Carrasco, E. *see* Hernández, C.
- Carrera, A. *see* Hernández, C.
- Carson, D.J. *see* Cardwell, C.R.
- Carter, A.M., Bennett, C.E., Bostock, J.A. & Grant, P.J. Metformin reduces C-reactive protein but not complement factor C3 in overweight patients with Type 2 diabetes mellitus, 1282
- Casamitjana, R. *see* Aguilera, E.
- Caselli, A., Latini, V., Lapenna, A., Di Carlo, S., Pirozzi, F., Benvenuto, A. & Uccioli, L. Transcutaneous oxygen tension monitoring after successful revascularization in diabetic patients with ischaemic foot ulcers, 460
- Cassader, M. *see* Bo, S.
- Cavuto, L. *see* Di Bonito, P.
- Cederholm, J. *see* Eliasson, B.
- Celermajer, D.S. *see* Lundman, P.
- Celermajer, D.S. *see* Keech, A.C.
- Ceriello, A. (Coordinator) for the SCAR (SCreening for ARteriopathy) study group *see* Faglia, E.
- Ceriello, A. PROactive Study: (r)evolution in the therapy of diabetes?, 1463
- Cha, D.R., Kim, I.S., Kang, Y.S., Han, S.Y., Han, K.H., Shin, C., Ji, Y.H. & Kim, N.H. Urinary concentration of transforming growth factor- $\beta$ -inducible gene-h3 ( $\beta$ ig-h3) in patients with Type 2 diabetes mellitus, 14
- Chan, A.K.F. *see* Donaghue, K.C.
- Chan, D.C. *see* Dogra, G.K.
- Channon, S. Huws-Thomas, M.V. Rollnick, S. & Gregory, J.W. The potential of motivational interviewing, 353
- Chantelau, E. The perils of procrastination: effects of early vs. delayed detection and treatment of incipient Charcot fracture, 1707
- Chapman, K.M. *see* Rutkove, S.B.
- Charbonnel, B.H., Matthews, D.R., Scherthaner, G., Hanefeld, M. & Brunetti, P. on behalf of the QUARTET Study Group. A long-term comparison of pioglitazone and gliclazide in patients with Type 2 diabetes mellitus: a randomized, double-blind, parallel-group comparison trial, 399
- Chat, S.W.S. *see* Choi, S.W.
- Chaturvedi, N. *see* Vèrier-Mine, O.
- Chatzianagnostou, K. *see* Di Cianni, G.
- Chaves, P. *see* Blaum, C.S.
- Cheema, F. *see* Maldonado, M.R.
- Chen, J.-W., Lauritzen, T., Christiansen, J.J., Jensen, L.H., Clausen, W.H.O. & Christiansen, J.S. Pharmacokinetic profiles of biphasic insulin aspart 30/70 and 70/30 in patients with Type 1 diabetes: a randomized double-blinded crossover study, 273
- Chen, L.-K., Chou, Y.-C., Tsai, S.-T., Hwang, S.-J. & Lee, S.-D. Hepatitis C virus infection-related Type 1 diabetes mellitus, 340
- Chen, S., Yan, W., Huang, J., Ge, D., Yao, Z. & Gu, D. Association analysis of the variant in the regulatory subunit of phosphoinositide 3-kinase ( $p85\alpha$ ) with Type 2 diabetes mellitus and hypertension in the Chinese Han population, 737
- Chen, Y.-C. *see* Hung, Y.-J.
- Chie, W.-C. *see* Lai, W.A.
- Chipchase, S. *see* Pound, N.
- Chipchase, S.Y., Treece, K.A., Pound, N., Game, F.L. & Jeffcoate, W.J. Heel ulcers don't heal in diabetes. Or do they? 1258
- Chirino, R. *see* Boronat, M.
- Chitson, P. *see* Söderberg, S.
- Chiu, K.C. *see* Jiang, Y.-D.
- Cho, N.-H. *see* Kim, D.-J.
- Cho, Y.M. *see* Lee, Y.Y.
- Cho, Y.M. *see* Moon, M.K.
- Choi, D.J. *see* Kwon, S.Y.
- Choi, S.W., Benzie, I.F.F., Lam, C.S.Y., Chat, S.W.S., Lam, J., Yiu, C.H., Kwan, J.J., Tang, Y.H., Yeung, G.S.P., Yeung, V.T.F., Woo, G.C., Hannigan, B.M. & Strain, J.J. Inter-relationships between DNA damage, ascorbic acid and glycaemic control in Type 2 diabetes mellitus, 1347
- Choi, Y.-H. *see* Kim, D.-J.
- Chokkalingam, K. *see* Kennedy, R.L.
- Chong, A.Y. *see* Lim, H.S.
- Chong, C.-K. *see* Tseng, C.-H.
- Chou, W.-L. *see* Jiang, Y.-D.
- Chou, Y.-C. *see* Chen, L.-K.
- Christiansen, J.J. *see* Chen, J.-W.
- Christiansen, J.S. *see* Chen, J.-W.
- Christiansen, J.S. *see* Sandbæk, A.
- Christie, M.R. *see* Damanhour, L.H.
- Chrysohoou, C. *see* Panagiotakos, D.B.
- Chrysohoou, C. *see* Pitsavos, C.
- Chuang, L.-M. *see* Jiang, Y.-D.
- Chung, M.G. *see* Kwon, S.Y.
- Church, C.O., Stevens, D.L. & Fugate, S.E. Diabetic ketoacidosis associated with aripiprazole, 1440
- Chwalow, A.J. *see* Du Pasquier-Fediaevsky, L.
- Ciardullo, A.V. Rosiglitazone improves insulin sensitivity, glucose tolerance and ambulatory blood pressure in subjects with impaired glucose tolerance: does it really work? If yes, is it a novelty? 666
- Ciccone, G. *see* Bo, S.
- Cicero, A.F.G., Dormi, A., Nascetti, S., Panourgia, M.P., Grandi, E., D'Addato, S. & Gaddi, A. Relative role of major risk factors for Type 2 diabetes development in the historical cohort of the Brisighella Heart Study: an 8-year follow-up, 1263
- Cilio, D. *see* Bruno, C.M.
- Clarey, A.J. *see* McMahon, S.K.
- Clark, P. *see* Middlebrooke, A.R.
- Clarke, G. *see* Wallymahmed, M.E.
- Clarke, N.R. & Narendran, P. Insulin prescribing is unsafe: education results in a significant but insufficient improvement, 1779
- Clarke, P., Norman, P., Coleman, M.A.G. & Holt, R. I.G. The introduction of a specific request form for the diagnosis of gestational diabetes (GDM) improves understanding of GDM amongst clinicians but does not increase its detection, 507
- Clausen, P. *see* Sørensen, V.R.
- Clausen, W.H.O. *see* Chen, J.-W.
- Cohen, R.M. *see* Schulze, M.B.
- Cohen, Y. *see* Wainstein, J.
- Colagiuri, S. *see* Glümer, C.
- Coleman, M.A.G. *see* Clarke, P.
- Coleman, R.L. *see* Stevens, R.J.
- Coll, A.P., Morganstein, D., Jayne, D., Soos, M.A., O'Rahilly, S. & Burke, J. Successful treatment of Type B insulin resistance in a

- patient with otherwise quiescent systemic lupus erythematosus, 814
- Colman, P.G. *see* Wraight, P.R.
- Conger, I. *see* Aguilera, E.
- Conne, P. *see* Pataky, Z.
- Conroy, C. *see* Simmons, D.
- Cook, D.G. *see* Whincup, P.H.
- Coppelli, A. *see* Giannarelli, R.
- Coresh, J. *see* Selvin, E.
- Corica, D. *see* Bonomo, M.
- Costa, L.A. *see* Canani, L.H.
- Coupaye, M. *see* Izzedine, H.
- Coupand, C. *see* Bebb, C.
- de Courten, M. *see* Söderberg, S.
- Courtney, C.H. *see* Au, S.
- Covino, G. *see* Di Bonito, P.
- Craig, M.E. *see* Donaghue, K.C.
- Crawford, M.A. *see* Min, Y.
- Creagh, N. Diabetic Foot—A Clinical Atlas, 816
- Crispim, D. *see* Canani, L.H.
- Crispim, D., Canani, L.H., Gross, J.L., Carlessi, R.M., Tschiedel, B., Souto, K.E.P. & Roisenberg, I. The G1888A variant in the mitochondrial 16S rRNA gene may be associated with Type 2 diabetes in Caucasian-Brazilian patients from southern Brazil, 1683
- Crock, P.A. *see* Donaghue, K.C.
- Csanády, M. *see* Nemes, A.
- Cuccuru, I. *see* Di Cianni, G.
- Cucinotta, D. *see* Perriello, G.
- Cull, C.A. *see* Thomason, M.J.
- Cummings, M.H. *see* Wotherspoon, F.
- Cundy, T. *see* Singh, R.
- Currie, C.J. *see* Lee, A.J.
- Curtis, J. *see* Wilson, C.
- Cusumano, J.M. *see* Donaghue, K.C.
- Czeizel, A.E. *see* Nielsen, G.L.
- D'Addato, S. *see* Cicero, A.F.G.
- D'Ambrogio, E., Giacomozzi, C., Macellari, V. & Uccioli, L. Abnormal foot function in diabetic patients: the altered onset of Windlass mechanism, 1713
- Dahlöf, B. *see* Hildebrandt, P.
- van Dam, R.M., Hoebee, B., Seidell, J.C., Schaap, M.M., de Bruin, T.W.A. & Feskens, E.J.M. Common variants in the ATP-sensitive K<sup>+</sup> channel genes *KCNJ11* (*Kir6.2*) and *ABCC8* (*SUR1*) in relation to glucose intolerance: population-based studies and meta-analyses, 590
- van Dam, R.M. *see* Pouwer, F.
- Damanhour, L.H., Dromey, J.A., Christie, M.R., Nasrat, H.A., Ardawi, M.S.M., Robins, R.A. & Todd, I. Autoantibodies to GAD and IA-2 in Saudi Arabian diabetic patients, 448
- Damsbo, P. *see* Hollingdal, M.
- Daneman, D. *see* Glaab, L.A.
- Daneman D. Is the 'Accelerator Hypothesis' worthy of our attention? 115
- Daniel, M. *see* Rowley, K.G.
- the Danish Study Group on Childhood Diabetes, *see* Nordly, S.
- Daousi, C., Benbow, S.J. & MacFarlane, I.A. Electrical spinal cord stimulation in the long-term treatment of chronic painful diabetic neuropathy, 393
- Datta, M. *see* Mohan, V.
- Davey Smith, G. *see* Lawlor, D.A.
- Davies, M.J. *see* Skinner, T.C.
- Davies, M.J. *see* Tringham, J.R.
- Davis, E.A. *see* McMahon, S.K.
- Davis, T.M.E., Bruce, D.G. & Davis, W.A. Predictors of first stroke in Type 1 diabetes: The Fremantle Diabetes Study, 551
- Davis, W.A. *see* Davis, T.M.E.
- Dawes, S. *see* Wallymahmed, M.E.
- Dawood, M. *see* George, P.M.
- De Berardis, G. *see* Franciosi, M.
- De Bellis, A. *see* Seghieri, G.
- Dean, H. Type 2 Diabetes in Childhood and Adolescence: A Global Perspective, 668
- The DECODE Study Group. Are insulin resistance, impaired fasting glucose and impaired glucose tolerance all equally strongly related to age? 1476
- Deepa, R. *see* Mohan, V.
- Deepa, R. *see* Vimalaswaran, K.S.
- Dekker, J.M. *see* Pouwer, F.
- Del Prato, S. *see* Di Cianni, G.
- Del Prato, S. *see* Giannarelli, R.
- Demers, L.M. *see* O'Bryan, T.A.
- Denniston, A.K.O., Banerjee, S., Gibson, J.M. & Dodson, P.M. Cardiovascular therapies and their role in diabetic eye disease, 665
- Deray, G. *see* Izzedine, H.
- Desha, Y.H. *see* Al-Bermani, A.
- Dethier, M. *see* Simoons, S.
- Devereux, R.B. *see* Hildebrandt, P.
- Dhatt, G.S. *see* Agarwal, M.M.
- Di Bonito, P., Moio, N., Cavuto, L., Covino, G., Murena, E., Scilla, C., Turco, S., Capaldo, B. & Sibilio, G. Early detection of diabetic cardiomyopathy: usefulness of tissue Doppler imaging, 1720
- Di Carlo, S. *see* Caselli, A.
- Di Cianni, G., Miccoli, R., Volpe, L., Lencioni, C., Ghio, A., Giovannitti, M.G., Cuccuru, I., Pellegrini, G., Chatzianagnostou, K., Boldrini, A. & Del Prato, S. Maternal triglyceride levels and newborn weight in pregnant women with normal glucose tolerance, 21
- Di Nardo, B. *see* Franciosi, M.
- Dimosthenopoulos, C. *see* Pitsavos, C.
- Dodson, P.M. *see* Denniston, A.K.O.
- Dogra, G.K., Watts, G.F., Chan, D.C. & Stanton, K. Statin therapy improves brachial artery vasodilator function in patients with Type 1 diabetes and microalbuminuria, 239
- Donaghue, K.C., Craig, M.E., Chan, A.K.F., Fairchild, J.M., Cusumano, J.M., Verge, C.F., Crock, P.A., Hing, S.J., Howard, N.J. & Silink, M. Prevalence of diabetes complications 6 years after diagnosis in an incident cohort of childhood diabetes, 711
- Donaghue, K.C., Margan, S.H., Chan, A.K.F., Holloway, B., Silink, M., Rangel, T. & Bennetts, B. The association of aldose reductase gene (*AKR1B1*) polymorphisms with diabetic neuropathy in adolescents, 1315
- Dong, Y., Gao, W., Nan, H., Yu, H., Li, F., Duan, W., Wang, Y., Sun, B., Qian, R., Tuomilehto, J. & Qiao, Q. Prevalence of Type 2 diabetes in urban and rural Chinese populations in Qingdao, China, 1427
- Donker, A.J.M. *see* Wulffelé, M.G.
- Donnan, P.T. *see* Donnelly, L.A.

- Donnelly, L.A., Morris, A.D., Frier, B.M., Ellis, J.D., Donnan, P.T., Durrant, R., Band, M.M., Reekie, G. & Leese, G.P. for the DARTS/MEMO Collaboration. Frequency and predictors of hypoglycaemia in Type 1 and insulin-treated Type 2 diabetes: a population-based study, 749
- Dormi, A. *see* Cicero, A.F.G.
- Dotevall, A., Hasdai, D., Wallentin, L., Battler, A. & Rosengren, A. Diabetes mellitus: clinical presentation and outcome in men and women with acute coronary syndromes. Data from the Euro Heart Survey ACS, 1542
- Douek, I.F., Allen, S.E., Ewings, P., Gale, E.A.M. & Bingley, P.J. for the Metformin Trial Group. Continuing metformin when starting insulin in patients with Type 2 diabetes: a double-blind randomized placebo-controlled trial, 634
- Douglas, M. *see* Lawton, J.
- Dowse, G.K. *see* Söderberg, S.
- Dozio, N. *see* Sampson, M.J.
- Draeger, E. *see* Pieber, T.R.
- Drivsholm, T. & de F. Olivarius, N. Routine diagnosis of Type 2 diabetes mellitus in general practice and hospitals: how do patients differ?, 336
- Drivsholm, T. *see* Rose, C.S.
- Dromey, J.A. *see* Damanhour, L.H.
- Du Pasquier-Fediaevsky, L., Chwalow, A.J., Tubiana-Rufi, N. & the PEDIAB Collaborative Group. Is the relationship between adherence behaviours and glycaemic control bi-directional at adolescence? A longitudinal cohort study, 427
- Duan, W. *see* Dong, Y.
- Duffy, N.A. *see* Lindsay, J.R.
- Dullaart, R.P.F. *see* Oomen, P.H.N.
- Durrant, R. *see* Donnelly, L.A.
- Eastwood, P.R. *see* Weisbrod, C.J.
- Ebrahim, S. *see* Lawlor, D.A.
- Ebrahim, S. *see* Thomas, M.C.
- Echwald, S. *see* Black, E.
- Edmonds, M., Foster, A. Wcer-free survival in diabetic foot patients 1293
- Edmonds, M. *see* Papanas, N.
- Edmonds, M.E. *see* Petrova, N.L.
- Edwards, R. *see* McElduff, P.
- Ekelund, U. *see* Särnblad, S.
- Elbagir, M. *see* Abdelgadir, M.
- Eliasson, B. *see* Fahlén, M.
- Eliasson, B., Cederholm, J., Nilsson, P. & Gudbjörnsdóttir, S. for the Steering Committee of the Swedish National Diabetes Register. The gap between guidelines and reality: Type 2 diabetes in a national diabetes register 1996–2003, 1420
- Elkmann, T. *see* Brassøe, R.
- Ellerbeck, E.F. *see* Bader, M.S.
- Ellingford, A. *see* Leese, G.P.
- Ellis, J.D. *see* Donnelly, L.A.
- Ellis, J.D. *see* Leese, G.P.
- Eltom, M. *see* Abdelgadir, M.
- Elving, L. *see* Bisseling, T.M.
- Embersson, J.R. *see* Thomas, M.C.
- Emslie-Smith, A. *see* McAlpine, R.R.
- Engelgau, M.M. *see* Simmons, D.
- Engelgau, M.M. *see* Tabaei, B.P.
- Ennis, C.N. *see* Au, S.
- Ercilla, G. *see* Aguilera, E.
- Eriksson, J.G. *see* Ylihäsälä, H.
- Ertugrul, D. *see* Sahin, M.
- Esnaola, S. *see* Larrañaga, I.
- Esteve, I. *see* Morcillo, S.
- The EURODIAB Prospective Complications Study Group. *See* Vèrier-Mine, O.
- Evans, E.M., Nye, F., Beeching, N.J. & Gill, G.V. 'Disappearing diabetes'—resolution of apparent Type 1 diabetes in a patient with AIDS and cytomegalovirus (CMV) infection, 218
- Evans, J.M.M. *see* McAlpine, R.R.
- Ewings, P. *see* Douek, I.F.
- Ewings, P. *see* Gooding, K.M.
- Fábregas, M. *see* Boronat, M.
- Faglia, E., Caravaggi, C., Marchetti, R., Mingardi, R., Morabito, A., Piaggese, A., Uccioli, L. & Ceriello, A. (Coordinator) for the SCAR (SCReening for ARterioopathy) study group Screening for peripheral arterial disease by means of the ankle-brachial index in newly diagnosed Type 2 diabetic patients, 1310
- Fahlén, M., Eliasson, B. & Odén, A. Optimization of basal insulin delivery in Type 1 diabetes: a retrospective study on the use of continuous subcutaneous insulin infusion and insulin glargine, 382
- Fairchild, J.M. *see* Donaghue, K.C.
- Falezza, G. *see* Targher, G.
- Fan, S.C. *see* Hung, Y.-J.
- Farmer, A. & Neil, A. In response to 'variations in glucose self-monitoring during oral hypoglycaemic therapy in primary care', 511
- Farmer, A., Gibson, O.J., Tarassenko, L. & Neil, A. A systematic review of telemedicine interventions to support blood glucose self-monitoring in diabetes, 1372
- Färnkqvist, L. *see* Åkesson, K.
- Farooqi, A.M. *see* Skinner, T.C.
- Farshchi, H.R. *see* Kennedy, R.L.
- Faure-Nogueras, E. *see* Gimeno-Orna, J.A.
- Fee, J.P.H. *see* McSorley, P.T.
- Fegan, P.G., Shore, A.C., Mawson, D., Tooke, J.E. & MacLeod, K.M. Microvascular endothelial function in subjects with Type 2 diabetes and the effect of lipid-lowering therapy, 1670
- Fehm, H.L. *see* Merl, V.
- Feinglos, M.N., Saad, M.F., Pi-Sunyer, F.X., An, B. & Santiago, O. on behalf of the Liraglutide Dose-Response Study Group, Effects of liraglutide (NN2211), a long-acting GLP-1 analogue, on glycaemic control and bodyweight in subjects with Type 2 diabetes, 1016
- Feldt-Rasmussen, B. *see* Sørensen, V.R.
- Felix Burden, A.C. Re: Maintaining glycaemic control during high-dose prednisolone administration in Type 1 diabetes, 510
- Fenger, M. *see* Bing, C.
- Fenger, M. *see* Koster, B.
- Fernández de la Puebla, R.A. *see* Paniagua, J.A.
- Fernandez-Real, J.M. *see* Vendrell, J.
- Ferrell, R.E. *see* Schneider, A.
- Feskens, E.J.M. *see* van Dam, R.M.
- the FIELD Vascular Study Investigators, *see* Keech, A.C.
- Fisher, L. *see* Armour, T.A.
- Fitzgerald, J.T. *see* Vijan, S.
- Flatman, M. *see* Sampson, M.J.



- Flatt, P.R. *see* Lindsay, J.R.  
 Flynn, A. *see* Perry, I.J.  
 Flyvbjerg, A. *see* Andersen, N.H.  
 Flyvbjerg, A. *see* Sørensen, V.R.  
 Fonollosa, A. *see* Hernández, C.  
 Forsblom, C. *see* Heinonen, P.  
 Forslund, L. *see* Held, C.  
 Foster, A. *see* Edmonds M. 1293  
 Forster, T. *see* Nemes, A.  
 Foster, A.V.M. *see* Petrova, N.L.  
 Foulon, E. *see* Simoens, S.  
 Fowkes, F.G.R. *see* Wild, S.H.  
 Fox, C. *see* Kilvert, A.  
 Frías, J.L. *see* Martínez-Frías, M.L.  
 Frías, J.P. *see* Martínez-Frías, M.L.  
 Földesi, I. *see* Bitó, T.  
 Franciosi, M., Peligrini, F., De Berardis, G., Belfiglio, M., Di Nardo, B., Greenfield, S., Kaplan, S.H., Rossi, M.C.E., Sacco, M., Tognoni, G., Valentini, M. & Nicolucci, A. for The QuED Study Group—quality of care and outcomes in Type 2 diabetes. Self-monitoring of blood glucose in non-insulin-treated diabetic patients: a longitudinal evaluation of its impact on metabolic control, 900  
 Francisco, G. *see* Hernández, C.  
 Franconi, F. *see* Seghieri, G.  
 Fraser, J. *see* Newton, D.J.  
 Fraser, R. Textbook of Diabetes and Pregnancy, 513  
 Frayling, T.M. *see* Zeggini, E.  
 Fredman, P. *see* Buschard, K.  
 Freestone, B. *see* Lim, H.S.  
 Frías, J.L. *see* Wyatt, J.W.  
 Fried, L.P. *see* Blaum, C.S.  
 Frier, B.M. *see* Donnelly, L.A.  
 Frier, B.M. *see* Graveling, A.J.  
 Frier, B.M. *see* Sommerfield, A.J.  
 Frier, B.M. *see* Strachan, M.W.J.  
 Froberg, K. *see* Lawlor, D.A.  
 Frölich, M. *see* Jazet, I.M.  
 Frommer-Shapira, R. *see* Rachmani, R.  
 Frossard, M., Blank, D., Joukhadar, Ch., Bayegan, K., Schmid, R., Luger, A. & Müller, M. Interstitial glucose in skeletal muscle of diabetic patients during an oral glucose tolerance test, 56  
 Fuentes, F. *see* Paniagua, J.A.  
 Fugate, S.E. *see* Church, C.O.  
 Fukui, M., Kitagawa, Y., Nakamura, N., Kadono, M., Hasegawa, G. & Yoshikawa, T. Association between alcohol consumption and serum dehydroepiandrosterone sulphate concentration in men with Type 2 diabetes: a link to decreased cardiovascular risk, 1446  
 Fuller, J.H. *see* Vèrier-Mine, O.  
 Furneri, S. *see* Perriello, G.  
 Furuta, H. *see* Nishi, M.  
 Gaddi, A. *see* Cicero, A.F.G.  
 Gale, E.A.M. *see* Douek, I.F.  
 Gale, E.A.M. & Atkinson, M. Reply, 958  
 Gall, M.-A. *see* Hollingdal, M.  
 Gallart, L. *see* Vendrell, J.  
 Gambino, R. *see* Bo, S.  
 Gamble, G. *see* Singh, R.  
 Game, F. *see* Idris, I.  
 Game, F. *see* Pound, N.  
 Game, F.L. *see* Chipchase, S.Y.  
 Gancia, R. *see* Bo, S.  
 Gao, W. *see* Dong, Y.  
 Gao, Y. *see* Shi, Y.-F.  
 Garceboo, H. *see* Söderberg, S.  
 Garg, S. *see* Wyatt, J.W.  
 Gatling, W. *see* Guzder, R.N.  
 Ge, D. *see* Chen, S.  
 Geiger, H. *see* Sanchez-Albisua, I.  
 Genschel, J. *see* Lüdtkke, A.  
 Gentile, L. *see* Bo, S.  
 George, P.M., Valabhji, J., Dawood, M. & Henry, J.A. Screening for Type 2 diabetes in the accident and emergency department, 1766  
 Gerds, E. *see* Hildebrandt, P.  
 Gershoni-Baruch, R. *see* Shehadeh, N.  
 Gfrerer, R.J. *see* Rakovac, I.  
 Ghebremeskel, K. *see* Min, Y.  
 Ghio, A. *see* Di Cianni, G.  
 Ghosh, S. *see* Vimalaswaran, K.S.  
 Giacomozzi, C. *see* D'Ambrogio, E.  
 Giannarelli, R., Coppelli, A., Boggi, U., Rizzo, G., Barsotti, M., Palaecologo, G., Tregnaighi, C., Mosca, F., Del Prato, S., Marchetti, P. New-onset diabetes after kidney transplantation, 1125  
 Gibson, J.M. *see* Denniston, A.K.O.  
 Gibson, O.J. *see* Farmer, A.  
 Gikas, A. *see* Sotiropoulos, A.  
 Giles, T. *see* Hildebrandt, P.  
 Gilg, J.A. *see* Whincup, P.H.  
 Gill, G. *see* Woodward, A.  
 Gill, G., Moulik, P. Mortality and diabetic neuropathy, 1289  
 Gill, G.V. *see* Evans, E.M.  
 Gill, G.V. *see* Patel, A.  
 Gill, G.V., Huddle, K.R.L. & Monkoe, G. Long-term (20 years) outcome and mortality of Type 1 diabetic patients in Soweto, South Africa, 1642  
 Gill, G.V., Morgan, C. & MacFarlane, I.A. Awareness and use of smoking cessation treatments among diabetic patients, 658  
 Gilliland, S. *see* Wilson, C.  
 Giménez-Pérez, G., Caixàs, A., Giménez-Palop, O., González-Clemente, J.M. & Mauricio, D. Dissemination of 'patient-oriented evidence that matters' on the Internet: the case of Type 2 diabetes treatment, 688  
 Giménez-Palop, O. *see* Giménez-Pérez, G.  
 Gimeno-Orna, J.A., Faure-Nogueras, E. & Sancho-Serrano, M.A. Usefulness of total cholesterol/HDL-cholesterol ratio in the management of diabetic dyslipidaemia, 26  
 Gin, H. *see* Rigalleau, V.  
 Giovannitti, M.G. *see* Di Cianni, G.  
 Glaab, L.A., Brown, R. & Daneman, D. School attendance in children with Type 1 diabetes, 421  
 Glaser, N.S. & Buonocore, M.H. Cerebral metabolic alterations in children with diabetic ketoacidosis, 515  
 Glümer, C. *see* Hamid, Y.H.  
 Glümer, C. *see* Rose, C.S.  
 Glümer, C., Borch-Johnsen, K. & Colagiuri, S. Can a screening programme for diabetes be applied to another population? 1234  
 Goh, L.-G. *see* Ng, T.-P.

- Gokulakrishnan, K. *see* Mohan, V.
- Golay, A. *see* Pataky, Z.
- Goldman, S.M. Nocturnal neuropathic pain in diabetic patients may be caused by spinal stenosis, 1763
- Gómez-Zumaquero, J.M. *see* Morcillo, S.
- Gomis, R. *see* Aguilera, E.
- Gonçalves dos Santos, K. *see* Canani, L.H.
- Gonçalves, D. *see* Bonomo, M.
- González, S., Kilpatrick, E.S. & Atkin, S.L. Improvement of glycaemic control with rebound following orlistat initiation and cessation associated with minimal weight change, 344
- González-Clemente, J.M. *see* Giménez-Pérez, G.
- Gonzalez, F. *see* Larrañaga, I.
- Gooding, K.M., MacLeod, K.M., Spyer, G., Ewings, P., Tooke, J.E. & Shore, A.C. Impact of hormone replacement therapy on microvascular function in healthy and Type 2 diabetic postmenopausal women, 536
- Grandi, E. *see* Cicero, A.F.G.
- Grant, P.J. *see* Carter, A.M.
- Grassi, G. *see* Bo, S.
- Graue, M., Wentzel-Larsen, T., Hanestad, B.R. & Søvik, O. Evaluation of a programme of group visits and computer-assisted consultations in the treatment of adolescents with Type 1 diabetes, 1522
- Graveling, A.J., Warren, R.E., Frier, B.M. Reply, 1290
- Gravholt, C.H. *see* Brassøe, R.
- Green, D.J. *see* Weisbrod, C.J.
- Greenfield, S. *see* Franciosi, M.
- Greenwood, R.H. *see* Sampson, M.J.
- Gregory, R. Unstable and Brittle Diabetes, 960
- Gregory, J.W. *see* Channon, S.
- Grieve, S.M. *see* Keech, A.C.
- Griffin, D. *see* Gunneberg, A.
- Griffiths, K. *see* Lundman, P.
- Griffiths, G. *see* Newton, D.J.
- Griffiths, K. *see* Keech, A.C.
- Grill, V. *see* Pieber, T.R.
- Grobbee, D.E. *see* Muis, M.J.
- Groshens, M. *see* Keech, A.C.
- Gross, J.L. *see* Canani, L.H.
- Gross, J.L. *see* Crispim, D.
- Gu, D. *see* Chen, S.
- Guang-da, X., Qiong-shu, W. & Wen, J. A T<sup>786</sup>C polymorphism in 5'-flanking region of the endothelial nitric oxide synthase gene and endothelium-dependent arterial dilation in Type 2 diabetes, 1663
- Gudbjörnsdóttir, S. for the Steering Committee of the Swedish National Diabetes Register *see* Eliasson, B.
- Guidi, S. *see* Bo, S.
- Gulliford, M. Cigarette smoking and prevention of Type 2 diabetes mellitus, 1285
- Gulliford, M.C. *see* Mahabir, D.
- Gunneberg, A. & Griffin, D. Laboratories should spot haemoglobin variants, 1779
- Guralnik, J.M. *see* Blaum, C.S.
- Gutiérrez, C. *see* Vendrell, J.
- Guvener, N.D. *see* Sahin, M.
- Guzder, R.N., Gatling, W., Mullee, M.A., Mehta, R.L. & Byrne, C.D. Prognostic value of the Framingham cardiovascular risk equation and the UKPDS risk equation for coronary heart disease in newly diagnosed Type 2 diabetes: results from a United Kingdom study, 554
- Haak, T. *see* Hermanns, N.
- Habacher, W. *see* Rakovac, I.
- Habeos, I.G., Theodoropoulou, A., Michalaki, M., Kyriazopoulou, V., Vagenakis, A.G., Psyrogiannis, A. Insulin sensitivity and fibrinogen concentrations in normoglycaemic offspring of Type 2 diabetic parents, 1122
- Hadley-Brown, M. *see* Sampson, M.J.
- Hagan Hughes, M. *see* Bosnyak, Z.
- Hagfelt, T. *see* Hoffmann, S.K.S.
- Halford, S. *see* Zeggini, E.
- Hamid, Y.H. *see* Rose, C.S.
- Hamid, Y.H., Vissing, H., Holst, B., Urhammer, S.A., Pyke, C., Hansen, S.K., Glümer, C., Borch-Johnsen, K., Jørgensen, T., Schwartz, T.W., Pedersen, O. & Hansen, T. Studies of relationships between variation of the human G protein-coupled receptor 40 Gene and Type 2 diabetes and insulin release, 74
- Hamman, R.F. *see* Schneider, A.
- Han, J-H. *see* Ko, S-H.
- Han, K.H. *see* Cha, D.R.
- Han, S.Y. *see* Cha, D.R.
- Hanefeld, M. *see* Charbonnel, B.H.
- Hanestad, B.R. *see* Graue, M.
- Hanley, A.J.G. *see* Pollex, R.L.
- Hannigan, B.M. *see* Choi, S.W.
- Hansen, K.W. *see* Andersen, N.H.
- Hansen, S.K. *see* Hamid, Y.H.
- Hansen, T. *see* Hamid, Y.H.
- Hansen, T. *see* Rose, C.S.
- Hansen, T. *see* Sagen, J.V.
- Hara, I. *see* Katahira, M.
- Harada, H. *see* Matsumura, Y.
- Harman-Boehm, I. *see* Wainstein, J.
- Harris, S., Ng, R., Syed, H. & Hillson, R. Near patient blood ketone measurements and their utility in predicting diabetic ketoacidosis, 221
- Harris, S.B. *see* Pollex, R.L.
- Harro, M. *see* Lawlor, D.A.
- Hartmann, R. *see* Kordonouri, O.
- Harvey, I. *see* Sampson, M.J.
- Harvey, J. Diabetes and the Kidney, 1127
- Hasdai, D. *see* Dotevall, A.
- Hasegawa, G. *see* Fukui, M.
- Hasegawa, K. *see* Yasuda, G.
- Hashimoto, K. *see* Ikeda, Y.
- Hattersley, A.T. *see* Sagen, J.V.
- Hattersley, A.T. *see* Zeggini, E.
- Hattori, H. *see* Oomen, P.H.N.
- Hayward, R.A. *see* Vijan, S.
- He, C.-T. *see* Hung, Y.-J.
- Heck, K. *see* Lüdtke, A.
- Hedner, J. *see* Buschard, K.
- Hegele, R.A. *see* Pollex, R.L.
- Heickendorff, L. *see* Andersen, N.H.
- Heine, R.J. *see* van der Ven, N.C.W.
- Heine, R.J. *see* Pouwer, F.
- Heinonen, P., Pettersson-Fernholm, K., Forsblom, C., Kallio, J., Scheinin, M. & P.-H. Groop and for the FinnDiane Study Group.

- A deletion polymorphism of the  $\alpha_{2B}$ -adrenergic receptor gene is not associated with late complications in Type 1 diabetic patients, 226
- Heise, M. *see* Bennett, S.M.A.
- Held, C., Björkander, I., Forslund, L., Rehnqvist, N. & Hjendahl, P. The impact of diabetes on elevated fasting blood glucose on cardiovascular prognosis in patients with stable angina pectoris, 1326
- Helleberg, K. *see* Andersen, N.H.
- Heller, R. *see* McElduff, P.
- Hemseloet, D.M. Acute brain injury with hemiparesis due to hypoglycaemia, 1287
- Hempel, M. *see* Brasso, R.
- Henareh, L., Jogestrand, T. & Agewall, S. Glucose intolerance is associated with C-reactive protein and intima-media anatomy of the common carotid artery in patients with coronary heart disease, 1212
- Henriksen, J.E. *see* Hoffmann, S.K.S.
- Henry, J.A. *see* George, P.M.
- Herman, W.H. *see* Tabaei, B.P.
- Hermann, N. *see* Nordly, S.
- Hermans, N., Kulzer, B., Krichbaum, M., Kubiak, T. & Haak, T. Affective and anxiety disorders in a German sample of diabetic patients: prevalence, comorbidity and risk factors, 293
- Hermans, M.P. *see* Buysschaert, M.
- Hernández, C., Lecube, A., Carrera, A. & Simó, R. Soluble transferrin receptors and ferritin in Type 2 diabetic patients, 97
- Hernández, C., Segura, R.M., Fonollosa, A., Carrasco, E., Francisco, G. & Simó, R. Interleukin-8, monocyte chemoattractant protein-1 and IL-10 in the vitreous fluid of patients with proliferative diabetic retinopathy, 719
- Heyburn, P.J. *see* Sampson, M.J.
- Higgins, C. *see* dos Santos Silva, I.
- Higgins, K.S. Diabetes and endocrinology as a career? A questionnaire survey of medical senior house officers, 955
- Hildebrandt, P., Wachtell, K., Dahlöf, B., Papadimitriou, V., Gerds, E., Giles, T., Oikarinen, L., Tuxen, C., Olsen, M.H. & Devereux, R.B. Impairment of cardiac function in hypertensive patients with Type 2 diabetes: a LIFE study, 1005
- Hill, J. *see* Shi, Y.-F.
- Hillson, R. *see* Harris, S.
- Hines, G. *see* Adler, A.I.
- Hing, S.J. *see* Donaghue, K.C.
- Hinthorn, D. *see* Bader, M.S.
- Hiramatsu, M. *see* Matsumura, Y.
- Hirota, Y. *see* Jimba, S.
- Hitman, G.A. *see* Boucher, B.J.
- Hitman, G.A. *see* Zeggini, E.
- Hjendahl, P. *see* Held, C.
- Hodis, H.N. *see* Karim, R.
- Hoebee, B. *see* van Dam, R.M.
- Hofer, T.P. *see* Vijan, S.
- Hoffmann, S.K.S., Jensen, L.O., Thyssen, P., Henriksen, J.E., Beck-Nielsen, H., Haghefelt, T. Patients with Type 2 diabetes and ischaemic heart disease have similar coronary flow reserve and endothelial function compared with non-diabetic patients with ischaemic heart disease, 1454
- Hoffstad, O. *see* Margolis, D.J.
- Hogenelst, M.H.E. *see* van der Ven, N.C.W.
- Hoi-Hansen, T., Pedersen-Bjergaard, U. & Thorsteinsson, B. Reproducibility and reliability of hypoglycaemic episodes recorded with Continuous Glucose Monitoring System (CGMS) in daily life, 858
- Hollingdal, M., Sturis, J., Gall, M.-A., Damsbo, P., Pincus, S., Veldhuis, J.D., Pøksen, N., Schmitz, O. & Juhl, C.B. Repaglinide treatment amplifies first-phase insulin secretion and high-frequency pulsatile insulin release in Type 2 diabetes, 1408
- Holloway, B. *see* Donaghue, K.C.
- Holman, R.R. *see* Adler, A.I.
- Holman, R.R. *see* Stevens, R.J.
- Holman, R.R. *see* Thomason, M.J.
- Holst, B. *see* Hamid, Y.H.
- Holst, C. *see* Black, E.
- Holst, J.J. *see* Hücking, K.
- Holt, R.I.G. *see* Clarke, P.
- Home, P.D., Home, E.M. International Textbook of Diabetes Mellitus, 1460
- Horio, T. *see* Miyazato, J.
- Howard, N.J. *see* Donaghue, K.C.
- Hoyme, H.E. *see* Wyatt, J.W.
- Hreidarsson, A.B. *see* Tryggvason, G.
- Hsieh, C.-H. *see* Hung, Y.-J.
- Hu, F.B. *see* Schulze, M.B.
- Huang, J. *see* Chen, S.
- Hubold, C. *see* Merl, V.
- Hücking, K., Kostic, Z., Pox, C., Ritzel, R., Holst, J.J., Schmiegel, W. & Nauck, M.A.  $\alpha$ -Glucosidase inhibition (acarbose) fails to enhance secretion of glucagon-like peptide 1 (7–36 amide) and to delay gastric emptying in Type 2 diabetic patients, 470
- Huddle, K.R.L. *see* Gill, G.V.
- Hung, Y.-J., Chen, Y.-C., Pei, D., Kuo, S.-W., Hsieh, C.-H., Wu, L.-Y., He, C.-T., Lee, C.H., Fan, S.C. & Sheu, W.H.-H. Sibutramine improves insulin sensitivity without alteration of serum adiponectin in obese subjects with Type 2 diabetes, 1024
- Hussain, A., Rahim, M.A., Azad Khan, A.K., Ali, S.M.K. & Vaaler, S. Type 2 diabetes in rural and urban population: diverse prevalence and associated risk factors in Bangladesh, 931
- Hutchison, A.S. Diabetes and Laboratory Medicine, 1127
- Huws-Thomas, M.V. *see* Channon, S.
- Hwang, S.-J. *see* Chen, L.-K.
- Ichiyanagi, K. *see* Yamanouchi, T.
- Idris, I., Game, F. & Jeffcoate, W. Does close glycaemic control promote healing in diabetic foot ulcers? Report of a feasibility study, 1060
- Igarashi, K. *see* Yamanouchi, T.
- Ikedo, Y., Suehiro, T., Osaki, F., Tsuzura, S., Arai, K., Kumon, Y. & Hashimoto, K. Interleukin-12B gene polymorphism is associated with circulating oxidized LDL levels in Japanese Type 2 diabetic patients, 661
- Ilonen, J. *see* Waris, M.E.
- Immonen-Räihä, P. *see* Pajunen, P.
- Inchiostro, S. Measurement of insulin sensitivity in Type 2 diabetes mellitus: comparison between  $K_{ITT}$  and HOMA-%S indices and evaluation of their resistance syndrome, 39
- Inder, T.E. *see* Cameron, F.J.
- Indran, M. *see* Kuppusamy, U.R.
- Indridason, O.S. *see* Tryggvason, G.

- Inenaga, T. *see* Miyazato, J.
- Inoue, Y. *see* Matsumura, Y.
- Ioannidis, I. *see* Pitsavos, C.
- the IONS study group, 803
- Ipp, E. *see* Linfoot, P.
- Ismail, K. *see* Rodrigues, I.A.S.
- Isogawa, A. *see* Waki, K.
- Ito, T. *see* Sakuta, H.
- Iwamoto, Y. *see* Jimba, S.
- Izawa, J. *see* Matsumura, Y.
- Izzedine, H., Coupaye, M., Reach, I. & Deray, G. Gastric bypass and resolution of proteinuria in an obese diabetic patient, 1761
- Jack Jr, L. *see* Armour, T.A.
- Jackson, S. *see* Kong M.-F.
- James, P. *see* McAlpine, R.R.
- Janatuinen, T. *see* Sundell, J.
- Järvi, A. *see* Abdelgadir, M.
- Jarvis, J. *see* Skinner, T.C.
- Jayne, D. *see* Brennan, A.M.
- Jayne, D. *see* Coll, A.P.
- Jazet, I.M., Pijl, H., Frölich, M., Schoemaker, R.C. & Meinders, A.E. Factors predicting the blood glucose lowering effect of a 30-day very low calorie diet in obese Type 2 diabetic patients, 52
- Jeffcoate, W. *see* Idris, I.
- Jeffcoate, W. *see* Pound, N.
- Jeffcoate, W.J. *see* Chipchase, S.Y.
- Jeitler, K. *see* Rakovac, I.
- Jensen, L.H. *see* Chen, J.-W.
- Jensen, L.O. *see* Hoffmann, S.K.S.
- Jerkkola, T. *see* Pajunen, P.
- Ji, Y.H. *see* Cha, D.R.
- Jiang, Y.-D., Yen, C.-J., Chou, W.-L., Kuo, S.-S., Lee, K.-C., Chiu, K.C. & Chuang, L.-M. Interaction of the G182C polymorphism in the APOA5 gene and fasting plasma glucose on plasma triglycerides in Type 2 diabetic subjects, 1690
- Jimba, S., Nakagami, T., Takahashi, M., Wakamatsu, T., Hirota, Y., Iwamoto, Y. & Wasada, T. Prevalence of non-alcoholic fatty liver disease and its association with impaired glucose metabolism in Japanese adults, 1141
- Jogestrand, T. *see* Henareh, L.
- Jogia, R. *see* Kong, M.-F.
- Johannesen, Ø. *see* Aas, A.M.
- Johansen, A. *see* Sagen, J.V.
- Johns, D. *see* Roden, M.
- Johnson, J.A., Simpson, S.H., Toth, E.L. & Majumdar, S.R. Reduced cardiovascular morbidity and mortality associated with metformin use in subjects with Type 2 diabetes, 497
- Jones, C. *see* Sampson, M.J.
- Jones, G. *see* McElduff, P.
- Jones, N.P. *see* Bennett, S.M.A.
- Jones, T.W. *see* McMahon, S.K.
- Jordahl, J. *see* Selvin, E.
- Jørgensen, T. *see* Bing, C.
- Jørgensen, T. *see* Hamid, Y.H.
- Jørgensen, T. *see* Nordly, S.
- Jørgensen, T. *see* Rose, C.S.
- Joukhadar, Ch. *see* Frossard, M.
- Jousilahti, P. *see* Yliharsilä, H.
- Jovanovic, L. *see* Wyatt, J.W.
- Juhl, C.B. *see* Hollingdal, M.
- Jung, H.S. *see* Lee, Y.Y.
- Jung, H.S. *see* Moon, M.K.
- Jung, J.-H. *see* Kim, D.-J.
- Jung, R.T. *see* Leese, G.P.
- Juolevi, A. *see* Pajunen, P.
- Kaaja, R. *see* Wyatt, J.W.
- Kadono, M. *see* Fukui, M.
- Kadowaki, T. *see* Waki, K.
- Kallio, J. *see* Heinonen, P.
- Kamide, K. *see* Miyazato, J.
- Kang, Y.S. *see* Cha, D.R.
- Kaplan, S.H. *see* Franciosi, M.
- Karges, B., Boehm, B.O. & Karges, W. Early hypoglycaemia after accidental intramuscular injection of insulin glargine, 1444
- Karges, W. *see* Karges, B.
- Kärjä-Koskenkari, P. *see* Pajunen, P.
- Karim, R., Buchanan, T.A., Hodis, H.N., Li, Y. & Mack, W.J. The association of smoking and subclinical atherosclerosis in Type 2 diabetes: modification by duration of diabetes, 81
- Karter, A.J., Ahmed, A.T., Liu, J., Moffet, H.H. & Parker, M.M. Pioglitazone initiation and subsequent hospitalization for congestive heart failure, 986
- Kasayama, S., Saito, H., Mukai, M. & Koga, M. Insulin sensitivity independently influences brachial-ankle pulse-wave velocity in non-diabetic subjects, 1701
- Katahira, M., Hara, I., Nishizaki, T. Insulin allergy decreased by Humulin S (Humulin R) and not by insulin aspart or Actrapid Penfill (Penfill R), 1455
- Katayama, Y. *see* Sakuta, H.
- Katsilambros, N. *see* Pitsavos, C.
- Kawai, M. *see* Nagaya, T.
- Kawano, Y. *see* Miyazato, J.
- Kawasaki, T. *see* Yamanouchi, T.
- Kean, M.J. *see* Cameron, F.J.
- Keech, A.C. *see* Lundman, P.
- Keech, A.C., Grieve, S.M., Patel, A., Griffiths, K., Skilton, M., Watts, G.F., Marwick, T.H., Groshens, M. & Celermajer, D.S., for the FIELD Vascular Study Investigators Urinary albumin levels in the normal range determine arterial wall thickness in adults with Type 2 diabetes: a FIELD substudy, 1558
- Kempler, P. *see* Nemes, A.
- Kendrick, D. *see* Bebb, C.
- Kendrick, D. on behalf of the Nottingham Diabetes Blood Pressure Study Group *see* Stewart, J.
- Kennedy, R.L., Chokkalingam, K. & Farshchi, H.R. Nutrition in patients with Type 2 diabetes: are low-carbohydrate diets effective, safe or desirable?, 821
- Kennedy-Martin, T. *see* Lee, A.J.
- Kern, W. *see* Merl, V.
- Kerr, L. *see* Wyatt, J.W.
- Kershaw, M.J.R., Newton, T., Barrett, T.G., Berry, K. & Kirk, J. Childhood diabetes presenting with hyperosmolar dehydration but without ketoacidosis: a report of three cases, 645
- Kesäniemi, Y.A. *see* Kunnari, A.
- Kesäniemi, Y.A. *see* Pajunen, P.
- Ketonen, M. *see* Pajunen, P.
- Khan, F. *see* Newton, D.J.
- Khan, H.M.R. *see* Pollex, R.L.

- Khanam, P.A. *see* Sayeed, M.A.  
 Khunti, K. *see* Skinner, T.C.  
 Khunti, K.K. *see* Tringham, J.R.  
 Kilpatrick, E.S. *see* González, S.  
 Kilvert, A. & Fox, C. Textbook of Type 2 Diabetes, 112  
 Kilvert, A. & Fox, C. Staged Diabetes Management, 2nd Edition, 961  
 Kim, D.-J., Noh, J.-H., Cho, N.-H., Lee, B.-W., Choi, Y.-H., Jung, J.-H., Min, Y.-K., Lee, M.-S., Lee, M.-K. & Kim, K.-W. Serum  $\gamma$ -glutamyltransferase within its normal concentration range is related to the presence of diabetes and cardiovascular risk factors, 1134  
 Kim, I.S. *see* Cha, D.R.  
 Kim, J.H. *see* Kwon, S.Y.  
 Kim, K.-W. *see* Kim, D.-J.  
 Kim, N.H. *see* Cha, D.R.  
 Kim, S.S. *see* Kwon, S.Y.  
 Kim, S.-R. *see* Ko, S.-H.  
 Kim, Y.K. *see* Kwon, S.Y.  
 Kim, Y.S. *see* Kwon, S.Y.  
 Kirk, J. *see* Kershaw, M.J.R.  
 Kisch, E. Diabetic patients with a HbA<sub>1c</sub> of 6.7% feel better than patients with a HbA<sub>1c</sub> of 9.1%, 1289  
 Kitagawa, Y. *see* Fukui, M.  
 Knip, M. *see* Waris, M.E.  
 Knudsen, S.T. *see* Andersen, N.H.  
 Knuuti, J. *see* Sundell, J.  
 Ko, S.-H., Lee, W.-Y., Lee, J.-H., Kwon, H.-S., Lee, J.-M., Kim, S.-R., Moon, S.-D., Song, K.-H., Han, J.-H., Ahn, Y.-B., Yoo, S.-J. & Son, H.-Y. Clinical characteristics of diabetic ketoacidosis in Korea over the past two decades, 466  
 Kockum, I. and the DISS study group *see* Åkesson, K.  
 Kocsis, B. *see* Szelestei, T.  
 Koga, M. *see* Kasayama, S.  
 Kogo, K. *see* Matsumura, Y.  
 Kohner, E.M. *see* Schneider, S.  
 Komiya, I. *see* Kouki, T.  
 Kong, M.-F., Jogia, R., Srinivasan, B.T. & Jackson, S. Malignant melanoma presenting as foot ulcer, 957  
 Kontela, E. *see* Sotiropoulos, A.  
 Koo, Y.S. *see* Kwon, S.Y.  
 Kooy, A. *see* Wulffélé, M.G.  
 Kordonouri, O., Hartmann, R. Higher body weight is associated with earlier onset of Type 1 diabetes in children: confirming the 'Accelerator Hypothesis', 1783  
 Koskinen, J.O. *see* Waris, M.E.  
 Koster, G. *see* Agarwal, M.M.  
 Køster B., Fenger M., Poulsen P., Vaag A. & Bentzen J. Novel polymorphisms in the GCKR gene and their influence on glucose and insulin levels in a Danish twin population, 1677  
 Kostic, Z. *see* Hücking, K.  
 Kouki, T., Takasu, N., Nakachi, A., Tamanaha, T., Komiya, I. & Tawata, M. Low-dose metformin improves hyperglycaemia related to myotonic dystrophy, 346  
 Koukkunen, H. *see* Pajunen, P.  
 Krichbaum, M. *see* Hermanns, N.  
 Kristensen, A. *see* Pieber, T.R.  
 Kubiak, T. *see* Hermanns, N.  
 Kuji, T. *see* Yasuda, G.  
 Kulzer, B. *see* Hermanns, N.  
 Kumakura, A. *see* Matsumura, Y.  
 Kumon, Y. *see* Ikeda, Y.  
 Kunnari, A., Ukkola, O. & Kesäniemi, Y.A. Resistin polymorphisms are associated with cerebrovascular disease in Finnish Type 2 diabetic patients, 583  
 Kuo, S.-S. *see* Jiang, Y.-D.  
 Kuo, S.-W. *see* Hung, Y.-J.  
 Kuppasamy, U.R., Indran, M. & Rokiah, P. Glycaemic control in relation to xanthine oxidase and antioxidant indices in Malaysian Type 2 diabetes patients, 1343  
 Kuulasmaa, K. *see* Pajunen, P.  
 Kwan, J.J. *see* Choi, S.W.  
 Kwon, H.-S. *see* Ko, S.-H.  
 Kwon, K.A. *see* Kwon, S.Y.  
 Kwon, O.S. *see* Kwon, S.Y.  
 Kwon, S.Y., Kim, S.S., Kwon, O.S., Kwon, K.A., Chung, M.G., Park, D.K., Kim, Y.S., Koo, Y.S., Kim, Y.K., Choi, D.J. & Kim, J.H. Prognostic significance of glycaemic control in patients with HBV and HCV-related cirrhosis and diabetes mellitus, 1530  
 Kyriazopoulou, V. *see* Habeos, I.G.  
 Laakso, M. *see* Roden, M.  
 Laekeman, G. *see* Simoons, S.  
 Lai, S.M. *see* Bader, M.S.  
 Lai, W.A., Lew-Ting, C.-Y. & Chie, W.-C. How diabetic patients think about and manage their illness in Taiwan, 286  
 Laine, H. *see* Sundell, J.  
 Laing, S.P. *see* dos Santos Silva, I.  
 Lam, C.S.Y. *see* Choi, S.W.  
 Lam, J. *see* Choi, S.W.  
 Lampasona, V. *see* Sordi, V.  
 Lanaras, L. *see* Sotiropoulos, A.  
 Lapenna, A. *see* Caselli, A.  
 Larrañaga, I., Arteagoitia, J.M., Rodriguez, J.L., Gonzalez, F., Esnaola, S. & Piniés, J.A. the Sentinel Practice Network of the Basque Country Socio-economic inequalities in the prevalence of Type 2 diabetes, cardiovascular risk factors and chronic diabetic complications in the Basque Country, Spain, 1047  
 Larrabee, J.E. *see* Rutkove, S.B.  
 Latini, V. *see* Caselli, A.  
 Lauritzen, T. *see* Chen, J.-W.  
 Lauritzen, T. *see* Peyrot, M.  
 Lauritzen, T. *see* Sandbæk, A.  
 Lawlor, D.A. *see* Thomas, M.C.  
 Lawlor, D.A., Ebrahim, S., Timpson, N. & Davey Smith, G. Avoiding milk is associated with a reduced risk of insulin resistance and the metabolic syndrome: findings from the British Women's Heart and Health Study, 808  
 Lawlor, D.A., Riddoch, C.J., Page, A.S., Anderssen, S.A., Froberg, K., Harro, M., Stansbie, D. & Davey Smith, G. The association of birthweight and contemporary size with insulin resistance among children from Estonia and Denmark: findings from the European Youth Heart Study, 921  
 Lawrence, E.C. *see* Schneider, A.  
 Lawrence, S.M. *see* Wraight, P.R.  
 Lawton, J., Parry, O., Peel, E. & Douglas, M. Diabetes service provision: a qualitative study of newly diagnosed Type 2 diabetes patients' experiences and views, 1246  
 Lecube, A. *see* Hernández, C.



- Lee, A.J. *see* Wild, S.H.
- Lee, A.J., Morgan, C.L.I., Morrissey, M., Wittrup-Jensen, K.U., Kennedy-Martin, T. & Currie, C.J. Evaluation of the association between the EQ-5D<sub>index</sub> (health-related utility) and body mass index (obesity) in hospital-treated people with Type 1 diabetes, Type 2 diabetes and with no diagnosed diabetes, 1482
- Lee, B.-W. *see* Kim, D.-J.
- Lee, C.H. *see* Hung, Y.-J.
- Lee, H.K. *see* Lee, Y.Y.
- Lee, H.K. *see* Moon, M.K.
- Lee, J.-H. *see* Ko, S.-H.
- Lee, J.-M. *see* Ko, S.-H.
- Lee, K. *see* Park, H.S.
- Lee, K.-C. *see* Jiang, Y.-D.
- Lee, M.-K. *see* Kim, D.-J.
- Lee, M.-S. *see* Kim, D.-J.
- Lee, N.S. *see* Lee, Y.Y.
- Lee, S.-D. *see* Chen, L.-K.
- Lee, W.-Y. *see* Ko, S.-H.
- Lee, Y.Y., Lee, N.S., Cho, Y.M., Moon, M.K., Jung, H.S., Park, Y.J., Park, H.J., Youn, B.S., Lee, H.K., Park, K.S. & Shin, H.D. Genetic association study of adiponectin polymorphisms with risk of Type 2 diabetes mellitus in Korean population, 569
- Lee-Parritz, A. *see* Wyatt, J.W.
- Leese, G.P. for the DARTS/MEMO Collaboration *see* Donnelly, L.A.
- Leese, G.P. *see* Newton, D.J.
- Leese, G.P., Morris, A.D., Swaminathan, K., Petrie, J.R., Sinharay, R., Ellingford, A., Taylor, A., Jung, R.T., Newton, R.W. & Ellis, J.D. Implementation of national diabetes retinal screening programme is associated with a lower proportion of patients referred to ophthalmology, 1112
- Lebert, P. *see* Wulffélé, M.G.
- Lehto, S. *see* Pajunen, P.
- Lehtonen, A. *see* Pajunen, P.
- Leinonen, P. *see* Loukovaara, M.
- Leiter, L.A.  $\beta$ -cell preservation: a potential role for thiazolidinediones to improve clinical care in Type 2 diabetes, 963
- Lencioni, C. *see* Di Cianni, G.
- Lengyel, C. *see* Nemes, A.
- Leong, H. *see* Ng, T.-P.
- Lernmark, Å. *see* Åkesson, K.
- Levy, J.C. *see* Zeggini, E.
- Levy, J.C. *see* Adler, A.I.
- Levy, Y. *see* Aronson, D.
- Lew-Ting, C.-Y. *see* Lai, W.A.
- Li, C.L. *see* Wang, X.L.
- Li, F. *see* Dong, Y.
- Li, J. *see* Olsen, J.
- Li, X. *see* Ran, X.
- Li, Y. *see* Karim, R.
- Lim, H.S. *see* Tan, K.T.
- Lim, H.S., Chong, A.Y., Freestone, B., Blann, A.D. & Lip, G.Y.H. The effect of multi-factorial intervention on plasma von Willebrand factor, soluble E-selectin and tissue factor in diabetes mellitus: implications for atherosclerotic vascular disease, 249
- Lindblad, U. *see* Buschard, K.
- Lindsay, J.R., Duffy, N.A., McKillop, A.M., Ardill, J., O'Harte, F.P.M., Flatt, P.R. & Bell, P.M. Inhibition of dipeptidyl peptidase IV activity by oral metformin in Type 2 diabetes, 654
- Lindström, J. *see* Yliharsilä, H.
- Linfoot, P., Bergstrom, C. & Ipp, E. Pathophysiology of ketoacidosis in Type 2 diabetes mellitus, 1414
- Lip, G.Y.H. *see* Lim, H.S.
- Lip, G.Y.H. *see* Tan, K.T.
- Lip, G.Y.H. *see* Tayebjee, M.H.
- Lipke, S. *see* Wilson, C.
- Lisbôa, H.R.K. *see* Canani, L.H.
- Liu, J. *see* Karter, A.J.
- Long, B. *see* McElduff, P.
- Lonovics, J. *see* Nemes, A.
- López-Miranda, J. *see* Paniagua, J.A.
- Loukovaara, M., Leinonen, P., Teramo, K., Nurminen, E., Andersson, S. & Rutanen, E.-M. Effect of maternal diabetes on phosphorylation of insulin-like growth factor binding protein-1 in cord serum, 434
- Lövestam-Adrian, M. *see* Torffvit, O.
- Lowy, C. *see* Min, Y.
- Lu, J.M. *see* Wang, X.L.
- Luger, A. *see* Frossard, M.
- Lüders, S., Schrader, J., Pfarr, E., Sturm, C.D., Paar, W.D. Strong association of C-reactive protein with 2-h-post-challenge glucose in non-diabetic subjects with hypertension, 1457
- Lüdtke, A., Heck, K., Genschel, J., Mehnert, H., Spuler, S., Worman, H.J. & Schmidt, H.H.-J. Long-term treatment experience in a subject with Dunnigan-type familial partial lipodystrophy: efficacy of rosiglitazone, 1611
- Lundman, P., Keech, A.C., Griffiths, K., Pillai, A. & Celermaier, D.S. New carotid plaque formation is very common in adult patients with Type 2 diabetes mellitus, 355
- Luo, Y. *see* Ran, X.
- Luotolahti, M. *see* Sundell, J.
- Lutale, J. *see* Abbas, Z.G.
- Luzio, S. *see* Schneider, S.
- Luzio, S.D. *see* Ollerton, R.L.
- Lympny, P.A. *see* Patel, A.
- Macellari, V. *see* D'Ambrogio, E.
- MacFadyen, R.J. *see* Tayebjee, M.H.
- MacFarlane, I.A. *see* Daousi, C.
- MacFarlane, I.A. *see* Gill, G.V.
- MacFarlane, I.A. *see* Wallymahmed, M.E.
- Mack, W.J. *see* Karim, R.
- MacLeod, K.M. *see* Fegan, P.G.
- MacLeod, K.M. *see* Gooding, K.M.
- MacLeod, K.M. *see* Middlebrooke, A.R.
- Madeley, R. *see* Bebb, C.
- Madsbad, S. *see* Bing, C.
- Mahabir, D. & Gulliford, M.C. Changing patterns of primary care for diabetes in Trinidad and Tobago over 10 years, 619
- Mähönen, M. *see* Pajunen, P.
- Mahtab, H. *see* Sayeed, M.A.
- Mai, K. *see* Sandbæk, A.
- Maia, A.L. *see* Canani, L.H.
- Maiello, M.R. *see* Spallone, V.
- Majumdar, S.R. *see* Johnson, J.A.
- Majumder, P.P. *see* Vimalaswaran, K.S.
- Makrilakis, K. *see* Pitsavos, C.
- Maldonado, M.R., Otiniano, M.E., Cheema, F., Rodriguez, L. & Balasubramanyam, A. Factors associated with insulin

- discontinuation in subjects with ketosis-prone diabetes but preserved  $\beta$ -cell function, 1744
- Malone, J.K., Bai, S., Campaigne, B.N., Reviriego, J. & Augendre-Ferrante, B. Twice-daily pre-mixed insulin rather than basal insulin therapy alone results in better overall glycaemic control in patients with Type 2 diabetes, 374
- Maltezos, E. *see* Papanas, N.
- Mann, R. *see* Sampson, M.J.
- Manson, J.E. *see* Schulze, M.B.
- Mannucci, E., Bardini, G. & Rotella, C. M. Effect of lower diagnostic thresholds on estimates of prevalence of impaired fasting glucose (IFG), 353
- Marin, C. *see* Paniagua, J.A.
- Marangou, D.A. *see* McMahon, S.K.
- Marchetti, P. *see* Giannarelli, R.
- Marchetti, R. *see* Faglia, E.
- Margan, S.H. *see* Donaghue, K.C.
- Margolis, D.J., Allen-Taylor, L., Hoffstad, O. & Berlin, J.A. Healing diabetic neuropathic foot ulcers: are we getting better? 172
- Mariz, S. *see* Roden, M.
- Marra, G. *see* Perriello, G.
- Marrero, D. *see* Boronati, M.
- Marshall, J.A. *see* Schneider, A.
- Marshall, S.M. Management of Diabetic Nephropathy, 668
- Marshall, S.M. Thoughts of an incoming Editor, 1
- Martínez-Frías, M.L., Frías, J.P., Bermejo, E., Rodríguez-Pinilla, E., Prieto, L. & Frías, J.L. Pre-gestational maternal body mass index predicts an increased risk of congenital malformations in infants of mothers with gestational diabetes, 775
- Marwick, T.H. *see* Keech, A.C.
- Mathiesen, E.R. *see* Sørensen, V.R.
- Mathieu, C. *see* Simoons, S.
- Matsumura, Y. *see* Waki, K.
- Matsumura, Y., Tsuji, A., Izawa, J., Kumakura, A., Hiramatsu, M., Kogo, K., Takahashi, K., Inoue, Y., Harada, H., Tonoizuka, N. Suspected toxicity from an iodoform preparation in a diabetic patient with multiple foot ulcers, 1121
- Matthews, D.R. *see* Adler, A.I.
- Matthews, D.R. *see* Charbonnel, B.H.
- Matthews, D.R. *see* Peyrot, M.
- Matthews, L.K. *see* Northam, E.A.
- Mattoo, V. *see* Wyatt, J.W.
- Mauricio, D. *see* Giménez-Pérez, G.
- Mawson, D. *see* Fegan, P.G.
- McAlpine, R.R., Morris, A.D., Emslie-Smith, A., James, P. & Evans, J.M.M. The annual incidence of diabetic complications in a population of patients with Type 1 and Type 2 diabetes, 348
- McCance, D.R. *see* McSorley, P.T.
- McCance, D.R. Diabetic Complications: New Diagnostic Tools and Therapeutic Advances, 669
- McCarthy, M.I. *see* Zeggini, E.
- McDermott, R. *see* Schutte, A.E.
- McDowell, J.R.S. *see* Moss, M.C.
- McElduff, P., Edwards, R., Burns, J.A., Young, R.J., Heller, R., Long, B., Jones, G. & New, J.P. Comparison of processes and intermediate outcomes between South Asian and European patients with diabetes in Blackburn, north-west England, 1226
- McElwee, K.J. *see* McMahon, S.K.
- McKillop, A.M. *see* Lindsay, J.R.
- McMahon, S.K., Airey, F.L., Marangou, D.A., McElwee, K.J., Carne, C.L., Clarey, A.J., Davis, E.A. & Jones, T.W. Insulin pump therapy in children and adolescents: improvements in key parameters of diabetes management including quality of life, 92
- McNamara, B. *see* Brennan, A.M.
- McSorley, P.T., Bell, P.M., Young, I.S., Atkinson, A.B., Sheridan, B., Fee, J.P.H. & McCance, D.R. Endothelial function, insulin action and cardiovascular risk factors in young healthy adult offspring of parents with Type 2 diabetes: effect of vitamin E in a randomized double-blind, controlled clinical trial, 703
- Meeking, D.R. *see* Wotherspoon, F.
- Megia, A. *see* Vendrell, J.
- Mehdi, M.Z. *see* Srivastava, A.K.
- Mehnert, H. *see* Lütke, A.
- Mehta, R.L. *see* Guzder, R.N.
- Meinders, A.E. *see* Jazet, I.M.
- Menzinger, G. *see* Spallone, V.
- Merati, R. *see* Bonomo, M.
- Merkouris, P. *see* Sotiropoulos, A.
- Merl, V., Peters, A., Oltmanns, K.M., Kern, W., Hubold, C., Fehm, H.L., Born, J. & Schultes, B. Antecedent hypoglycaemia attenuates vascular endothelial growth factor response to subsequent hypoglycaemia in healthy men, 1278
- Merletti, F. *see* Bo, S.
- Metelko, Z. *see* Pibernik-Okanovic, M.
- Metzger, M. *see* Wainstein, J.
- Meyer, C., Tolia, A., Platanisiotis, D., Stumvoll, M., Vlachos, L. & Mitrakou, A. Increased renal glucose metabolism in Type 1 diabetes mellitus, 453
- Miccoli, R. *see* Di Cianni, G.
- Michalaki, M. *see* Habeos, I.G.
- Middlebrooke, A.R., Armstrong, N., Welsman, J.R., Shore, A.C., Clark, P. & MacLeod, K.M. Does aerobic fitness influence microvascular function in healthy adults at risk of developing Type 2 diabetes?, 483
- Miettinen, H. *see* Pajunen, P.
- Min, Y., Lowy, C., Ghebremeskel, K., Thomas, B., Bitsanis, D. & Crawford, M.A. Fetal erythrocyte membrane lipids modification: preliminary observation of an early sign of compromised insulin sensitivity in offspring of gestational diabetic women, 914
- Min, Y.-K. *see* Kim, D.-J.
- Mingardi, R. *see* Faglia, E.
- Minuchin, O. *see* Wainstein, J.
- Mion, E. *see* Bonomo, M.
- Missoni, E. *see* Mlinaric Missoni, E.
- Mitrakou, A. *see* Meyer, C.
- Mittleman, M.A. *see* Rana, J.S.
- Miyazato, J., Horio, T., Takiuchi, S., Kamide, K., Sasaki, O., Nakamura, S., Nakahama, H., Inenaga, T., Takishita, S. & Kawano, Y. Left ventricular diastolic dysfunction in patients with chronic renal failure: impact of diabetes mellitus, 730
- Mlinaric Missoni, E., Vukelic, M., de Soy, D., Belicza, M., Vazic Babic, V., Missoni, E. Fungal infection in diabetic foot ulcers, 1124
- Moffet, H.H. *see* Karter, A.J.
- Mogensen, C.E. *see* Andersen, N.H.
- Mohan, V. *see* Adaikalakoteswari, A.
- Mohan, V. *see* Vimalaswaran, K.S.
- Mohan, V., Deepa, R., Velmurugan, K. & Premalatha, G. Association of C-reactive protein with body fat, diabetes and

- coronary artery disease in Asian Indians: The Chennai Urban Rural Epidemiology Study (CURES-6), 863
- Mohan, V., Gokulakrishnan, K., Deepa, R., Shanthirani, C.S. & Datta, M. Association of physical inactivity with components of metabolic syndrome and coronary artery disease—the Chennai Urban Population Study (CUPS no. 15), 1206
- Moio, N. *see* Di Bonito, P.
- Mojiminiyi, O.A. *see* Abdella, N.A.
- Molnar, G.A. *see* Szelestei, T.
- Molven, A. *see* Sagen, J.V.
- Monden, T. *see* Shimizu, H.
- Monkoe, G. *see* Gill, G.V.
- Moon, M.K. *see* Lee, Y.Y.
- Moon, M.K., Cho, Y.M., Jung, H.S., Park, Y.J., Yoon, K.H., Sung, Y.A., Park, B.L., Lee, H.K., Park, K.S. & Shin, H.D. Genetic polymorphisms in peroxisome proliferator-activated receptor  $\gamma$  are associated with Type 2 diabetes mellitus and obesity in the Korean population, 1161
- Moon, S-D. *see* Ko, S-H.
- Morabito, A. *see* Bonomo, M.
- Morabito, A. *see* Faglia, E.
- Morcillo, S., Cardona, F., Rojo-Martínez, G., Esteve, I., Ruiz-de-Adana, M.S., Tinañones, F., Gómez-Zumaquero, J.M. & Soriguer, F. Association between MspI polymorphism of the APO AI gene and Type 2 diabetes mellitus, 782
- Morgan, C. *see* Gill, G.V.
- Morgan, C.Ll. *see* Lee, A.J.
- Morgan, J. *see* Al-Bermani, A.
- Morgan, J.P. *see* Rana, J.S.
- Morganstein, D. *see* Brennan, A.M.
- Morganstein, D. *see* Coll A.P.
- Mori, M. *see* Shimizu, H.
- Morris, A.D. *see* Donnelly, L.A.
- Morris, A.D. *see* Leese, G.P.
- Morris, A.D. *see* McAlpine, R.R.
- Morris, A.D. *see* dos Santos Silva, I.S.
- Morrissey, M. *see* Lee, A.J.
- Mortality and diabetic neuropathy, 1289
- Mortensen, H.B. *see* Nordly, S.
- Mosca, F. *see* Giannarelli, R.
- Moss, M.C. & McDowell, J.R.S. Rural Vincentians' (Caribbean) beliefs about the usage of non-prescribable medicines for treating Type 2 diabetes, 1492
- Motta, G. *see* Bonomo, M.
- Moulik, P. *see* Gill, G.
- Moussa, M.A. *see* Abdella, N.A.
- Mrak, P. *see* Rakovac, I.
- Muis, M.J., Bots, M.L., Grobbee, D.E. & Stolk, R.P. Insulin treatment and cardiovascular disease; friend or foe? A point of view, 118
- Mukai, M. *see* Kasayama, S.
- Mukamal, K.J. *see* Rana, J.S.
- Mulle, M.A. *see* Gunder, R.N.
- Müller, M. *see* Frossard, M.
- Muller, J.E. *see* Rana, J.S.
- Mulligan, J. *see* Betts, P.
- Munday, L.J. *see* Wotherspoon, F.
- Murena, E. *see* Di Bonito, P.
- Murphy, H.R. *see* Roland, J.M.
- Mustonen, J. *see* Pajunen, P.
- Nagai, T. *see* Shimizu, H.
- Nagaya, T., Yoshida, H., Takahashi, H. & Kawai, M. Increases in body mass index, even within non-obese levels, raise the risk for Type 2 diabetes mellitus: a follow-up study in a Japanese population, 1107
- Nagy, I. *see* Nemes, A.
- Nagy, J. *see* Szelestei, T.
- Naides, S.J. *see* O'Bryan, T.A.
- Nakachi, A. *see* Kouki, T.
- Nakagami, T. *see* Jimba, S.
- Nakahama, H. *see* Miyazato, J.
- Nakamura, N. *see* Fukui, M.
- Nakamura, S. *see* Miyazato, J.
- Nakao, T. *see* Nishi, M.
- Nan, H. *see* Dong, Y.
- Nanjo, K. *see* Nishi, M.
- Narendran, P. *see* Clarke, N.R.
- Nascetti, S. *see* Cicero, A.F.G.
- Nasrat, H.A. *see* Damanhour, L.H.
- Nauck, M.A. *see* Hücking, K.
- Naum, A. *see* Sundell, J.
- Neil, A. *see* Farmer, A.
- Neil, A. *see* Stratton, I.M.
- Neil, J.J. *see* Cameron, F.J.
- Nekolla, S.G. *see* Bengel, F.M.
- Nemes, A., Lengyel, C., Forster, T., Várkonyi, T.T., Takács, R., Nagy, I., Kempler, P., Lonovics, J. & Csanády, M. Coronary flow reserve, insulin resistance and blood pressure response to standing in patients with normoglycaemia: is there a relationship?, 1614
- Nester, C. *see* Thompson, L.
- Nesto, R.W. *see* Rana, J.S.
- Neu, A. *see* Sanchez-Albisua, I.
- Nerve, J. *see* Bengel, F.M.
- New, J.P. *see* McElduff, P.
- Newton, D.J., Bennett, S.P., Fraser, J., Khan, F., Belch, J.J.F., Griffiths, G. & Leese, G.P. Pilot study of the effects of local pressure on microvascular function in the diabetic foot, 1487
- Newton, R.W. *see* Leese, G.P.
- Newton, T. *see* Kershaw, M.J.R.
- Ng, R. *see* Harris, S.
- Ng, T.-P., Goh, L.-G., Tan, Y., Tan, E., Leong, H., Tay, E.-G. & Thai, A.-C. Ethnic differences in glycaemic control in adult Type 2 diabetic patients in primary care: a 3-year follow-up study, 1598
- N'Guyen Thi, N., Paries, J., Attali, J.R. & Valensi, P. High prevalence and severity of cardiac autonomic neuropathy in Vietnamese diabetic patients, 1072
- Nicoletti, F. *see* Aguilera, E.
- Nicolucci, A. for The QuED Study Group—quality of care and outcomes in Type 2 diabetes *see* Franciosi, M.
- Nielsen, G.L., Norgard, B., Puho, E., Rothman, K.J., Sorensen, H.T. & Czeizel, A.E. Risk of specific congenital abnormalities in offspring of women with diabetes, 693
- Nielsen, S. *see* Andersen, H.
- Niemelä, M. *see* Pajunen, P.
- Nijpels, G. *see* Pouwer, F.
- Nikolaou, T. *see* Sotiropoulos, A.
- Nilsson, P. *see* Eliasson, B.
- Nishi, M., Sasahara, M., Shono, T., Saika, S., Yamamoto, Y., Ohkawa, K., Furuta, H., Nakao, T., Sasaki, H. & Nanjo, K. A

- case of novel de novo paired box gene 6 (PAX6) mutation with early-onset diabetes mellitus and aniridia, 641
- Nishimura, R. *see* Bosnyak, Z.
- Nishizaki, T. *see* Katahira, M.
- Njølstad, P.R. *see* Sagen, J.V.
- Njølstad, P.R. *see* Shehadeh, N.
- Noda, M. *see* Waki, K.
- Noh, J.-H. *see* Kim, D.-J.
- Nordly, S., Mortensen, H.B., Andreassen, A.H., Hermann, N. & Jørgensen, T., on behalf of the Danish Study Group on Childhood Diabetes, Factors associated with glycaemic outcome of childhood diabetes care in Denmark, 1566
- Norman, P. *see* Clarke, P.
- Nørgård, B. *see* Nielsen, G.L.
- Norris, J.M. *see* Schneider, A.
- Norris, S.L. *see* Armour, T.A.
- Northam, E.A., Matthews, L.K., Anderson, P.J., Cameron, F.J. & Werther, G.A. Psychiatric morbidity and health outcome in Type 1 diabetes – perspectives from a prospective longitudinal study, 152
- Northcote-Wright, J. *see* Roland, J.M.
- the Nottingham Diabetes Blood Pressure Study Group, 152
- Nóvoa, F.J. *see* Boronat, M.
- Nurminen, E. *see* Loukovaara, M.
- Nuutila, P. *see* Sundell, J.
- Nyári, T. *see* Bitó, T.
- Nye, F. *see* Evans, E.M.
- Nyström, L. *see* Åkesson, K.
- O'Halloran, D.J. *see* Brennan, A.M.
- O'Harte, F.P.M. *see* Lindsay, J.R.
- O'Bryan, T.A., Beck, M.J., Demers, L.M., Naides, S.J. Human parvovirus B19 infection in children with new onset Type 1 diabetes mellitus, 1778
- Odén, A. *see* Fahlén, M.
- O'Dea, K. *see* Rowley, K.G.
- O'Dea, K. *see* Schutte, A.E.
- Odoki, K. *see* Whincup, P.H.
- O'Driscoll, G. *see* Weisbrod, C.J.
- Ogawa, N. *see* Yasuda, G.
- Ohashi, Y. *see* Waki, K.
- Ohkawa, K. *see* Nishi, M.
- Ohshiro, Y., Takasu, N. Reply, 1288
- Oikarinen, L. *see* Hildebrandt, P.
- Okoro, E.O. *see* Ologe, F.E.
- de F. Olivarius, N. *see* Drivsholm, T.
- Ollerton, R.L., Luzio, S.D. & Owens, D.R. Contribution of fasting and postprandial plasma glucose to HbA<sub>1c</sub>, 954
- Ologe, F.E. & Okoro, E.O. Type 2 diabetes and hearing loss in black Africans, 664
- Olsen, J., Li, J. & Precht, D.H. Hospitalization because of diabetes and bereavement: a national cohort study of parents who lost a child, 1338
- Olsen, M.H. *see* Hildebrandt, P.
- Oltmanns, K.M. *see* Merl, V.
- Oomen, P.H.N., van Tol, A., Hattori, H., Smit, A.J., Scheek, L.M. & Dullaart, R.P.F. Human plasma phospholipid transfer protein activity is decreased by acute hyperglycaemia: studies without and with hyperinsulinaemia in Type 1 diabetes mellitus, 768
- O'Rahilly, S. *see* Coll, A.P.
- Orchard, T.J. *see* Bosnyak, Z.
- Oriola, J. *see* Aguilera, E.
- Orr, N.J. & Boyages, S.C. Patterns of care as a risk factor for the development of vision-threatening diabetic retinopathy: a population-based matched case-control study using insurance claims (Medicare) data, 1083
- Osaki, F. *see* Ikeda, Y.
- Östman, J. *see* Åkesson, K.
- Oswald, G. & Yudkin, J.S. Prognostic value of admission plasma glucose and HbA<sub>1c</sub> in acute myocardial infarction, 509
- Otiniano, M.E. *see* Maldonado, M.R.
- Owen, C.G. *see* Whincup, P.H.
- Owen, K.R. *see* Zeggini, E.
- Owens, D.R. *see* Ollerton, R.L.
- Owens, D.R. *see* Schneider, S.
- Ozer, H., Selcuk, D., Albayram, S. Acute brain injury in hypoglycaemia-induced hemiplegia, 1287
- Paar, W.D. *see* Lüders, S.
- Pagano, G.F. *see* Bo, S.
- Page, A.S. *see* Lawlor, D.A.
- Page, S. Emergencies in diabetes diagnosis, management and prevention, 816
- Pajunen, P., Koukkunen, H., Ketonen, M., Jerkkola, T., Immonen-Räihä, P., Kärjä-Koskenkari, P., Mähönen, M., Niemelä, M., Kuulasmaa, K., Palomäki, P., Mustonen, J., Lehtonen, A., Arstila, M., Vuorenmaa, T., Lehto, S., Miettinen, H., Juolevi, A., Torppa, J., Tuomilehto, J., Kesäniemi, Y.A., Pyörälä, K. & Salomaa, V. Five-year risk of developing clinical diabetes after first myocardial infarction; the FINAMI study, 1334
- Pál, A. *see* Bitó, T.
- Palaeologo, G. *see* Giannarelli, R.
- Palmer, A.J. *see* Roze, S.
- Palomäki, P. *see* Pajunen, P.
- Palsson, R. *see* Tryggvason, G.
- Pampanelli, S. *see* Perriello, G.
- Pan, C.Y. *see* Wang, X.L.
- Pan, C.-Y. *see* Shi, Y.-F.
- Panagiotakos, D.B. *see* Pitsavos, C.
- Panagiotakos, D.B., Pitsavos, C., Chrysohou, C. & Stefanadis, C. The epidemiology of Type 2 diabetes mellitus in Greek adults: the ATTICA study, 1581
- Paniagua, J.A., López-Miranda, J., Pérez-Martínez, P., Marín, C., Vida, J.M., Fuentes, F., Fernández de la Puebla, R.A. & Pérez-Jiménez, F. Oxidized-LDL levels are changed during short-term serum glucose variations and lowered with statin treatment in early Type 2 diabetes: a study of endothelial function and microalbuminuria, 1647
- Panourgia, M.P. *see* Cicero, A.F.G.
- Papademitriou, V. *see* Hildebrandt, P.
- Papanas, N., Edmonds, M. & Maltezos, E. Pseudoclaudication as a manifestation of diabetic neuropathy, 1608
- Pappas, S. *see* Sotiropoulos, A.
- Paries, J. *see* N'Guyen Thi, N.
- Park, B.L. *see* Moon, M.K.
- Park, D.K. *see* Kwon, S.Y.
- Park, H.J. *see* Lee, Y.Y.
- Park, H.S. & Lee, K. Greater beneficial effects of visceral fat reduction compared with subcutaneous fat reduction on

- parameters of the metabolic syndrome: a study of weight reduction programmes in subjects with visceral and subcutaneous obesity, 266
- Park, K.S. *see* Lee, Y.Y.
- Park, K.S. *see* Moon, M.K.
- Park, Y.J. *see* Lee, Y.Y.
- Park, Y.J. *see* Moon, M.K.
- Parker, L. *see* Adams, J.
- Parker, M.M. *see* Karter, A.J.
- Parkinson, J.R.C. *see* Zeggini, E.
- Parry, O. *see* Lawton, J.
- Pataký, Z., Assal, J.-P., Conne, P., Vuagnat, H. & Golay, A. Plantar pressure distribution in Type 2 diabetic patients without peripheral neuropathy and peripheral vascular disease, 762
- Patel, A. *see* Keech, A.C.
- Patel, A., Scott, W.R., Lympany, P.A., Rippin, J.D., Gill, G.V., the Warren 3/UK GoKind Study Group, Barnett, A.H. & Bain, S.C. The TGF- $\beta$ 1 gene codon 10 polymorphism contributes to the genetic predisposition to nephropathy in Type 1 diabetes, 69
- Patterson, C.C. *see* Cardwell, C.R.
- Pearce, M.S. *see* Adams, J.
- Pearson, D.W.M. *see* dos Santos Silva, I.
- Pearson, E.R. *see* Sagen, J.V.
- Pedersen, O. *see* Black, E.
- Pedersen, O. *see* Hamid, Y.H.
- Pedersen, O. *see* Rose, C.S.
- Pedersen, O. *see* Sagen, J.V.
- Pedersen-Bjergaard, U. *see* Hoi-Hansen, T.
- the PEDIAB Collaborative Group, 427
- Peel, E. *see* Lawton, J.
- Pei, D. *see* Hung, Y.-J.
- Pellegrini, F. *see* Franciosi, M.
- Pellegrini, G. *see* Di Cianni, G.
- Pellicano, R. *see* Bruno, C.M.
- Pellicano, R. *see* Clarke, N.R.
- Peppas, T. *see* Sotiropoulos, A.
- Pérez-Jiménez, F. *see* Paniagua, J.A.
- Pérez-Martínez, P. *see* Paniagua, J.A.
- Perlemoine, C. *see* Rigalleau, V.
- Peros, K. *see* Pibernik-Okanovic, M.
- Perriello, G., Pampanelli, S., Porcellati, F., Avogaro, A., Bosi, E., Petrella, G., Squatrito, S., Furneri, S., Marra, G., Vitali, L., Previti, M. & Cucinotta, D. Insulin aspart improves meal time glycaemic control in patients with Type 2 diabetes: a randomized, stratified, double-blind and cross-over trial, 606
- Perry, I.J., Villegas, R., Salim, A. & Flynn, A. Clustering of protective factors for glucose intolerance and insulin resistance: a cross-sectional study, 1091
- Peters, A. *see* Merl, V.
- Petersson, G.H. *see* Twetman, S.
- Petrella, G. *see* Perriello, G.
- Petrie, J.R. *see* Leese, G.P.
- Petrova, N.L., Foster, A.V.M. & Edmonds, M.E. Calcaneal bone mineral density in patients with Charcot neuropathic osteoarthropathy: differences between Type 1 and Type 2 diabetes, 756
- Pettersson-Fernholm, K. *see* Heinonen, P.
- Peyrot, M., Rubin, R.R., Lauritzen, T., Snoek, F.J., Matthews, D.R. & Skovlund, S.E. on behalf of the International DAWN Advisory Panel Psychosocial problems and barriers to improved diabetes management: results of the Cross-National Diabetes Attitudes, Wishes and Needs (DAWN) Study, 1379
- Pfarr, E. *see* Lüders, S.
- P.-H. Group and for the FinnDiane Study Group, 226
- Piaggese, A. *see* Faglia, E.
- Pibernik-Okanovic, M., Peros, K., Szabo, S., Begic, D. & Metelko, Z. Depression in Croatian Type 2 diabetic patients: prevalence and risk factors. A Croatian survey from the European Depression in Diabetes (EDID) Research Consortium, 942
- Pieber, T.R. for the FQSD Austria *see* Rakovac, I.
- Pieber, T.R., Draeger, E., Kristensen, A. & Grill, V. Comparison of three multiple injection regimens for Type 1 diabetes: morning plus dinner or bedtime administration of insulin detemir vs. morning plus bedtime NPH insulin, 850
- Pijl, H. *see* Jazet, I.M.
- Pillai, A. *see* Lundman, P.
- Pincus, S. *see* Hollingdal, M.
- Piniés, J.A. *see* Larrañaga, I.
- Pirozzi, F. *see* Caselli, A.
- Pi-Sunyer, F.X. *see* Feinglos, M.N.
- Pitsavos, C. *see* Panagiotakos, D.B.
- Pitsavos, C., Makrilakis, K., Panagiotakos, D.B., Chrysoshoou, C., Ioannidis, I., Dimosthenopoulos, C., Stefanadis, C. & Katsilambros, N. The J-shape effect of alcohol intake on the risk of developing acute coronary syndromes in diabetic subjects: the CARDIO2000 II Study, 243
- Plant, W.D. *see* Brennan, A.M.
- Platanisiotis, D. *see* Meyer, C.
- van der Ploeg, H.M. *see* van der Ven, N.C.W.
- Pollex, R.L., Hanley, A.J.G., Zinman, B., Harris, S.B., Khan, H.M.R. & Hegele, R.A. Synergism between mutant *HNFI1A* and the metabolic syndrome in Oji-Cree Type 2 diabetes, 1510
- Porcellati, F. *see* Perriello, G.
- Pørksen, N. *see* Hollingdal, M.
- Poulsen, P. *see* Køster, B.
- Poulsen, P.L. *see* Andersen, N.H.
- Poulsen, S.H. *see* Andersen, N.H.
- Pound, N. *see* Chipchase, S.Y.
- Pound, N., Chipchase, S., Treece, K., Game, F. & Jeffcoate, W. Ulcer-free survival following management of foot ulcers in diabetes, 1306
- Pouwer, F., Nijpels, G., Beekman, A.T., Dekker, J.M., van Dam, R.M., Heine, R.J. & Snoek, F.J. Fat food for a bad mood. Could we treat and prevent depression in Type 2 diabetes by means of  $\omega$ -3 polyunsaturated fatty acids? A review of the evidence, 1465
- Pox, C. *see* Hücking, K.
- Precht, D.H. *see* Olsen, J.
- Premalatha, G. *see* Mohan, V.
- Previti, M. *see* Perriello, G.
- Prieto, L. *see* Martínez-Frías, M.L.
- Prognostic value of admission plasma glucose and HbA<sub>1c</sub> in acute myocardial infarction, 509
- Promoting targeted screening for Type 2 diabetes mellitus: the contribution of community pharmacists, 812
- Psyrogiannis, A. *see* Habeos, I.G.
- Puho, E. *see* Nielsen, G.L.
- Punnose, J. *see* Agarwal, M.M.
- Pyrälä, K. *see* Pajunen, P.
- Pyke, C. *see* Hamid, Y.H.



- Qian, R. *see* Dong, Y.  
 Qiao, Q. *see* Dong, Y.  
 Qiong-shu, W. *see* Guang-da, X.
- Rachmani, R., Slavachevski, I., Berla, M., Frommer-Shapira, R. & Ravid, M. Teaching and motivating patients to control their risk factors retards progression of cardiovascular as well as microvascular sequelae of Type 2 diabetes mellitus—a randomized prospective 8 years follow-up study, 410  
 Radha, V. *see* Vimalaswaran, K.S.  
 Ragusa, A. *see* Bonomo, M.  
 Rahim, M.A. *see* Hussain, A.  
 Rahu, M. *see* Ainla, T.  
 Raitakari, O.T. *see* Sundell, J.  
 Rakovac, I., Jeitler, K., Gfrerer, R.J., Habacher, W., Seereiner, S., Mrak, P. & Pieber, T.R. for the FQSD Austria. Patients with Type 2 diabetes treated with metformin: prevalence of contraindications and their correlation with discontinuation, 662  
 Ran, X., Wang, C., Wang, H., Zhao, T., Tong, N., Song, B., Bu, H., Luo, Y., Tian, H. & Li, X. Muscle infarction involving muscles of abdominal and thoracic walls in diabetes, 1757  
 Rana, J.S., Mukamal, K.J., Nesto, R.W., Morgan, J.P., Muller, J.E. & Mittleman, M.A. Effect of diabetes mellitus and its treatment on ventricular arrhythmias complicating acute myocardial infarction, 576  
 Rangel, T. *see* Donaghue, K.C.  
 Rao, M.R.S. *see* Vimalaswaran, K.S.  
 Råstam, L. *see* Buschard, K.  
 Ravid, M. *see* Rachmani, R.  
 Raz, I. *see* Wainstein, J.  
 Reach, G. Role of habit in adherence to medical treatment, 415  
 Reach, I. *see* Izzedine, H.  
 Reekie, G. *see* Donnelly, L.A.  
 Rees, D.A. & Alcolado, J.C. Animal models of diabetes mellitus, 359  
 Rehnqvist, N. *see* Held, C.  
 Reid, H.A. *see* Rodrigues, I.A.S.  
 Reviriego, J. *see* Malone, J.K.  
 Ricart, W. *see* Vendrell, J.  
 Richards, R.J. Regarding the use of HOMA to aid drug selection, 225  
 Richardson, C. *see* Roden, M.  
 Richart, C. *see* Vendrell, J.  
 Riddoch, C.J. *see* Lawlor, D.A.  
 Rifai, N. *see* Schulze, M.B.  
 Rigalleau, V., Baillet-Blanco, L., Perlemoine, C., Salmi, L.-R. & Gin, H. Lower plasma triglycerides are associated with increased need for insulin requirement in poorly controlled Type 2 diabetic patients, 877  
 Rippin, J.D. *see* Patel, A.  
 Ritzel, R. *see* Hücking, K.  
 Rizzo, G. *see* Giannarelli, R.  
 Robins, R.A. *see* Damanhour, L.H.  
 Robinson, A. *see* Robinson, S.  
 Robinson, S., Robinson, A. Hypoglycaemia and driving, 1289  
 Roden, M., Laakso, M., Johns, D., Widel, M., Urquhart, R., Richardson, C., Mariz, S. & Tan, M.H. Long-term effects of pioglitazone and metformin on insulin sensitivity in patients with Type 2 diabetes mellitus, 1101  
 Rodríguez-Pinilla, E. *see* Martínez-Frías, M.L.  
 Rodrigues, I.A.S., Reid, H.A., Ismail, K. & Amiel, S.A. Indications and efficacy of continuous subcutaneous insulin infusion (CSII) therapy in Type 1 diabetes mellitus: a clinical audit in a specialist service, 842  
 Rodriguez, J.L. *see* Larrañaga, I.  
 Rodriguez, L. *see* Maldonado, M.R.  
 Roisenberg, I. *see* Canani, L.H.  
 Roisenberg, I. *see* Crispim, D.  
 Rojo-Martínez, G. *see* Morcillo, S.  
 Rokiah, P. *see* Kuppusamy, U.R.  
 Roland, J.M., Murphy, H.R., Ball, V., Northcote-Wright, J. & Temple, R.C. The pregnancies of women with Type 2 diabetes: poor outcomes but opportunities for improvement, 1774  
 Rollnick, S. *see* Channon, S.  
 Ronis, D.L. *see* Vijan, S.  
 Rönnemaa, T. *see* Sundell, J.  
 Rosato, R. *see* Bo, S.  
 Rose, C.S., Andersen, G., Hamid, Y.H., Glümer, C., Drivsholm, T., Borch-Johnsen, K., Jørgensen, T., Pedersen, O. & Hansen, T. Studies of relationships between the GLUT10 Ala206Thr polymorphism and impaired insulin secretion, 946  
 Rosengren, A. *see* Dotevall, A.  
 Rossi, M.C.E. *see* Franciosi, M.  
 Rotella, C.M. *see* Mannu, E.  
 Rothman, K.J. *see* Nielsen, G.L.  
 Rowley, K. *see* Schutte, A.E.  
 Rowley, K.G., Daniel, M. & O'Dea, K. Screening for diabetes in Indigenous populations using glycated haemoglobin: sensitivity, specificity, post-test likelihood and risk of disease, 833  
 Royal College of Physicians of Edinburgh Diabetes Register Group. Predicting vascular risk in Type 1 diabetes: stratification in a hospital based population in Scotland, 164  
 Royle, P.L., Bain, L. & Waugh, N.R. Sources of evidence for systematic reviews of interventions in diabetes, 1386  
 Roze, S., Valentine, W.J., Zakrzewska, K.E. & Palmer, A.J. Health-economic comparison of continuous subcutaneous insulin infusion with multiple daily injection for the treatment of Type 1 diabetes in the UK, 1239  
 Ruiz-de-Adana, M.S. *see* Morcillo, S.  
 Rubin, R.R. *see* Peyrot, M.  
 Russel, F.G.M. *see* Bisseling, T.M.  
 Rutanen, E.-M. *see* Loukovaara, M.  
 Rutkove, S.B., Chapman, K.M., Acosta, J.A. & Larrabee, J.E. Foot temperature in diabetic polyneuropathy: innocent bystander or unrecognized accomplice? 231  
 Rutten, G. Diabetes patient education: time for a new era, 671  
 Saad, M.F. *see* Feinglos, M.N.  
 Saavedra, P. *see* Boronat, M.  
 Sacco, M. *see* Franciosi, M.  
 Sagen, J.V., Baumann, M.E., Salvesen, H.B., Molven, A., Søvik, O. & Njølstad, P.R. Diagnostic screening of *NEURODI* (MODY6) in subjects with MODY or gestational diabetes mellitus, 1012  
 Sagen, J.V., Pearson, E.R., Johansen, A., Spyer, G., Søvik, O., Pedersen, O., Njølstad, P.R., Hattersley, A.T. & Hansen, T. Preserved insulin response to tolbutamide in hepatocyte nuclear factor-1 $\alpha$  mutation carriers, 406

- Sahin, M., Bakiner, O., Ertugrul, D., Guvener, N.D. Creatine kinase elevation in a patient taking rosiglitazone, 1624
- Saika, S. *see* Nishi, M.
- Saito, H. *see* Kasayama, S.
- Sakai, T. *see* Yamanouchi, T.
- Sakuta, H., Suzuki, T., Katayama, Y., Yasuda, H. & Ito, T. Heavy alcohol intake, homocysteine and Type 2 diabetes, 1359
- Salim, A. *see* Perry, I.J.
- Salmi, L.-R. *see* Rigalleau, V.
- Salomaa, V. *see* Pajunen, P.
- Salvesen, H.B. *see* Sagen, J.V.
- Sampson, M.J. *see* Zeggini, E.
- Sampson, M.J., Barrie, P., Dozio, N., Flatman, M., Hadley-Brown, M., Harvey, I., Heyburn, P.J., Jones, C., Mann, R., Temple, R.C. & Greenwood, R.H. A mobile screening programme for the cardiovascular and microvascular complications of Type 2 diabetes in primary care, 256
- Sanchez-Albisua, I., Wolf, J., Neu, A., Geiger, H., Wäscher, I. & Stern, M. Coeliac disease in children with Type 1 diabetes mellitus: the effect of the gluten-free diet, 1079
- Sancho-Serrano, M.A. *see* Gimeno-Orna, J.A.
- Sandbæk, A., Lauritzen, T., Borch-Johnsen, K., Mai, K. & Christiansen, J.S. The comparison of venous plasma glucose and whole blood capillary glucose in diagnoses of Type 2 diabetes: a population-based screening study, 1173
- dos Santos Silva, I., Higgins, C., Swerdlow, A.J., Laing, S.P., Slater, S.D., Pearson, D.W.M. & Morris, A.D. Birthweight and other pregnancy outcomes in a cohort of women with pre-gestational insulin-treated diabetes mellitus, Scotland, 1979-95, 440
- Särnblad, S., Ekelund, U. & Åman, J. Physical activity and energy intake in adolescent girls with Type 1 diabetes, 893
- Sarturi Tres, G. *see* Canani, L.H.
- Sasahara, M. *see* Nishi, M.
- Sasaki, H. *see* Nishi, M.
- Sasaki, O. on behalf of the liraglutide Dose-Response Study Group *see* Miyazato, J.
- Sasaki, S. *see* Waki, K.
- Sato, T. *see* Shimizu, H.
- Saunders, S. *see* Wallymahmed, M.E.
- Sayeed, M.A., Mahtab, H., Khanam, P.A., Begum, R., Banu, A. & Azad Khan, A.K. Diabetes and hypertension in pregnancy in a rural community of Bangladesh: a population-based study, 1267
- Scala, L. *see* Targher, G.
- Scarpello, J.H.B. *see* Buch, H.N.
- Schaap, M.M. *see* van Dam, R.M.
- Scheek, L.M. *see* Oomen, P.H.N.
- Scheinin, M., *see* Heinonen, P.
- Scherbaum, W.A. *see* Schinner, S.
- Schernthaner, G. *see* Belcher, G.
- Schernthaner, G. *see* Charbonnel, B.H.
- Schinner, S., Scherbaum, W.A., Bornstein, S.R. & Barthel, A. Molecular mechanisms of insulin resistance, 674
- Schmid, R. *see* Frossard, M.
- Schmidt, H.H.-J. *see* Lüdtk, A.
- Schmidt, V. *see* Schneider, S.
- Schmieg, W. *see* Hüking, K.
- Schmitz, O. *see* Andersen, H.
- Schmitz, O. *see* Hollingdal, M.
- Schneider, A., Lawrence, E.C., Barmada, M.M., Norris, J.M., Hamman, R.F., Marshall, J.A., Ferrell, R.E. & Whitcomb, D.C. The SPINK1 N34S mutation is not associated with Type 2 diabetes mellitus in a population of the USA, 744
- Schneider, S., Aldington, S.J., Kohner, E.M., Luzio, S., Owens, D.R., Schmidt, V., Schuell, H. & Zahlmann, G. Quality assurance for diabetic retinopathy telecreening, 794
- Schnell, O. *see* Bengel, F.M.
- Schoemaker, R.C. *see* Jazet, I.M.
- Schrader, J. *see* Lüders, S.
- Schuell, H. *see* Schneider, S.
- Schultes, B. *see* Merl, V.
- Schulze, M.B., Solomon, C.G., Rifai, N., Cohen, R.M., Sparrow, J., Hu, F.B. & Manson, J.E. Hyperproinsulinaemia and risk of Type 2 diabetes mellitus in women, 1178
- Schutte, A.E., Shemesh, T., Rowley, K., Best, J.D., McDermott, R. & O'Dea, K. The metabolic syndrome and changing relationship between blood pressure and insulin with age, as observed in Aboriginal and Torres Strait Islander peoples, 1589
- Schwaiger, M. *see* Bengel, F.M.
- Schwartz, T.W. *see* Hamid, Y.H.
- Sciaccia, C. *see* Bruna, C.M.
- Scilla, C. *see* Di Bonito, P.
- Scott, W.R. *see* Patel, A.
- Seely, E.W. *see* Wyatt, J.W.
- Seereiner, S. *see* Rakovac, I.
- Seghieri, G., De Bellis, A., Anichini, R., Alviggi, L., Franconi, F. & Breschi, M.C. Does parity increase insulin resistance during pregnancy?, 1574
- Segura, R.M. *see* Hernández, C.
- Seidell, J.C. *see* van Dam, R.M.
- Selcuk, D. *see* Ozer, H.
- Selvin, E., Coresh, J., Jordahl, J., Boland, L. & Steffes, M.W. Stability of haemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) measurements from frozen whole blood samples stored for over a decade, 1726
- the Sentinel Practice Network of the Basque Country, 1047
- Shanthirani, C.S. *see* Mohan, V.
- Shaw, J.E. *see* Söderberg, S.
- Shaw, K.M. *see* Wotherspoon, F.
- Shehadeh, N., Bakri, D., Njølstad, P.R. & Gershoni-Baruch, R. Clinical characteristics of mutation carriers in a large family with glucokinase diabetes (MODY2), 994
- Shemesh, T. *see* Schutte, A.E.
- Sheridan, B. *see* Au, S.
- Sheridan, B. *see* McSorley, P.T.
- Sheu, J.-J. *see* Tseng, C.-H.
- Sheu, W.H.-H. *see* Hung, Y.-J.
- Shi, Y.-F., Pan, C.-Y., Hill, J. & Gao, Y. Orlistat in the treatment of overweight or obese Chinese patients with newly diagnosed Type 2 diabetes, 1737
- Shimizu, H., Monden, T., Nagai, T., Shoda, Y., Sato, T., Yamada, M. & Mori, M. Insulin resistance determines efficacy of glimepiride in Type 2 diabetic patients not well controlled by diet alone, 225
- Shimura, G. *see* Yasuda, G.
- Shin, C. *see* Cha, D.R.
- Shin, H.D. *see* Lee, Y.Y.
- Shin, H.D. *see* Moon, M.K.
- Shoda, Y. *see* Shimizu, H.

- Shono, T. *see* Nishi, M.
- Shore, A.C. *see* Fegan, P.G.
- Shore, A.C. *see* Gooding, K.M.
- Shore, A.C. *see* Middlebrooke, A.R.
- Sibilio, G. *see* Di Bonito, P.
- Silink, M. *see* Donaghue, K.C.
- Silli, B. *see* Bo, S.
- Simó, R. *see* Hernández, C.
- Simón, I. *see* Vendrell, J.
- Simell, O. *see* Waris, M.E.
- Simmons, D., Conroy, C. & Thompson, C.F. In-hospital breast feeding rates among women with gestational diabetes and pregestational Type 2 diabetes in South Auckland, 177
- Simmons, D., Thompson, C.F. & Englgau, M.M. Controlling the diabetes epidemic: how should we screen for undiagnosed diabetes and dysglycaemia? 207
- Simoens, S., Foulon, E., Dethier, M., Mathieu, C. & Laekeman, G. Promoting targeted screening for Type 2 diabetes mellitus: the contribution of community pharmacists, 812
- Simpson, S.H. *see* Johnson, J.A.
- Singh, R., Gamble, G. & Cundy, T. Lifetime risk of symptomatic carpal tunnel syndrome in Type 1 diabetes, 625
- Sinharay, R. *see* Leese, G.P.
- Skilton, M. *see* Keech, A.C.
- Skinner, T.C., Davies, M.J., Farooqi, A.M., Jarvis, J., Tringham, J.R. & Khunti, K. Diabetes screening anxiety and beliefs, 1497
- Skliros, E. *see* Sotiropoulos, A.
- Skourtis, S. *see* Sotiropoulos, A.
- Skovlund, S.E. on behalf of the International DAWN Advisory Panel *see* Peyrot, M.
- Slater, S. *see* Vijan, S.
- Slater, S.D. *see* dos Santos Silva, I.
- Slavachevski, I. *see* Rachmani, R.
- Smit, A.J. *see* Oomen, P.H.N.
- Smith, B. *see* Betts, P.
- Smith, F.B. *see* Wild, S.H.
- Smits, P. *see* Bisseling, T.M.
- Smits, P. *see* Veldman, B.A.J.
- Snoek, F.J. *see* van der Ven, N.C.W.
- Snoek, F.J. *see* Peyrot, M.
- Snoek, F.J. *see* Pouwer, F.
- Söderberg, S., Zimmet, P., Tuomilehto, J., de Courten, M., Dowse, G.K., Chitson, P., Gareeboo, H., Alberti, K.G.M.M. & Shaw, J.E. Increasing prevalence of Type 2 diabetes mellitus in all ethnic groups in Mauritius, 61
- Soós, A. *see* Barkai, L.
- Solberg, M. *see* Aas, A.M.
- Solomon, C.G. *see* Schulze, M.B.
- Sommerfield, A.J. & Frier, B.M. A new cause of neuroglycopenia: 'missing the point'—impaired visual function during acute hypoglycaemia, 510
- Son, H.-Y. *see* Ko, S.-H.
- Song, B. *see* Ran, X.
- Song, K.-H. *see* Ko, S.-H.
- Soobrah, R. *see* Al-Bermani, A.
- Soos, M.A. *see* Brennan, A.M.
- Soos, M.A. *see* Coll, A.P.
- Sordi, V., Lampasona, V., Cainarca, S. & Bonifacio, E. No evidence of diabetes-specific CD38 (ADP ribosyl cyclase/cyclic ADP-ribose hydrolase) autoantibodies by liquid-phase immunoprecipitation, 1770
- Sørensen, H.T. *see* Nielsen, G.L.
- Sørensen, T.I.A. *see* Black, E.
- Sørensen, V.R., Mathiesen, E.R., Clausen, P., Flyvbjerg, A. & Feldt-Rasmussen, B. Impaired vascular function during short-term poor glycaemic control in Type 1 diabetic patients, 871
- Soriguer, F. *see* Morcillo, S.
- Sotiropoulos, A., Merkouris, P., Gikas, A., Skourtis, S., Skliros, E., Lanaras, L., Nikolaou, T. & Pappas, S. Influenza and pneumococcal vaccination rates among Greek diabetic patients in primary care, 110
- Sotiropoulos, A., Skourtis, S., Merkouris, P., Peppas, T., Apostolou, O., Kontela, E., Skliros, E., Pappas, S. Incidence and outcome of asymptomatic bacteriuria in females with Type 2 diabetes mellitus over a 1-year follow-up period and association with risk factors, 1625
- Souto, K.E.P. *see* Crispim, D.
- Søvik, O. *see* Graue, M.
- Søvik, O. *see* Sagen, J.V.
- de Soy, D. *see* Mlinaric Missoni, E.
- Spallone, V., Maiello, M.R. & Menzinger G. Reply, 1781
- Sparrow, J. *see* Schulze, M.B.
- Spuler, S. *see* Lüdtkke, A.
- Spyer, G. *see* Gooding, K.M.
- Spyer, G. *see* Sagen, J.V.
- Squatraro, S. *see* Perriello, G.
- Srinivasan, B.T. *see* Kong, M.-F.
- Srivastava, A.K. & Mehdi, M.Z. Insulino-mimetic and anti-diabetic effects of vanadium compounds, 2
- Standl, E. *see* Bengel, F.M.
- Stansbie, D. *see* Lawlor, D.A.
- Stanton, K. *see* Dogra, G.K.
- Steevers, E.A.P. *see* Bisseling, T.M.
- Stefanadis, C. *see* Panagiotakos, D.B.
- Stefanadis, C. *see* Pitsavos, C.
- Steffes, M.W. *see* Selvin, E.
- Stehouwer, C.D.A. *see* Wulffélé, M.G.
- Stern, M. *see* Sanchez-Albisua, I.
- Stevens, D.L. *see* Church, C.O.
- Stevens, R.J., Coleman, R.L. & Holman, R.R. Framingham risk equations underestimate coronary heart disease risk in diabetes, 228
- Stewart, J. & Kendrick, D. on behalf of the Nottingham Diabetes Blood pressure Study Group. Setting and negotiating blood sugar and blood pressure targets with people with Type 2 diabetes in primary care: a cross sectional survey, 683
- Stewart, J. *see* Bebb, C.
- Stolk, R.P. *see* Muis, M.J.
- Strachan, M.W.J. & Frier, B.M. Cognitive impairment following severe hypoglycaemia: response to Kubiak *et al.*, 959
- Strain, J.J. *see* Choi, S.W.
- Stratton, I.M. & Neil, A. How to ensure your paper is rejected by the statistical reviewer, 371
- Stratton, I.M. *see* Adler, A.I.
- Stuart, L. *see* Thompson, L.
- Stuart, N.S. *see* Vijan, S.
- Stumvoll, M. *see* Meyer, C.
- Sturis, J. *see* Hollingdal, M.
- Sturm, C.D. *see* Lüders, S.

- Sturrock, N. for the Nottingham Diabetes Blood Pressure Study *see* Bebb, C.
- Suehiro, T. *see* Ikeda, Y.
- Sun, B. *see* Dong, Y.
- Sundell, J., Janatuinen, T., Rönnemaa, T., Naum, A., Laine, H., Luotolahti, M., Nuutila, P., Raitakari, O.T. & Knuuti, J. Lifetime glycaemic exposure predicts reduced coronary vasoreactivity in Type 1 diabetic subjects, 45
- Sundvall, J. *see* Ylihäsälä, H.
- Sung, Y.A. *see* Moon, M.K.
- Suzuki, T. *see* Sakuta, H.
- Swaminathan, K. *see* Leese, G.P.
- Swerdlow, A.J. *see* dos Santos Silva, I.
- Syed, H. *see* Harris, S.
- Symonds, C.S. *see* Al-Bermani, A.
- Szabo, S. *see* Pibernik-Okanovic, M.
- Szelestei, T., Bähring, S., Wagner, Z., Aydin, A., Molnár, G.A., Kocsis, B., Nagy, J., Wittmann, I. Serum levels of L-arginine analogues and glutathione peroxidase and catalase gene variants in Type 2 diabetes mellitus patients, 356
- Tabaei, B.P., Engelgau, M.M. & Herman, W.H. A multivariate logistic regression equation to screen for dysglycaemia: development and validation, 599
- Tajima, N. *see* Bosnyak, Z.
- Takács, R. *see* Nemes, A.
- Takahashi, H. *see* Nagaya, T.
- Takahashi, K. *see* Matsumura, Y.
- Takahashi, M. *see* Jimba, S.
- Takahashi, Y. *see* Waki, K.
- Takasu, N. *see* Kouki, T.
- Takasu, N. *see* Ohshiro, Y.
- Takishita, S. *see* Miyazato, J.
- Takiuchi, S. *see* Miyazato, J.
- Tamanaha, T. *see* Kouki, T.
- Tan, E. *see* Ng, T.-P.
- Tan, K.T., Tayebjee, M.H., Lim, H.S. & Lip, G.Y.H. Clinically apparent atherosclerotic disease in diabetes is associated with an increase in platelet microparticle levels, 1657
- Tan, M. *see* Wyatt, J.W.
- Tan, M.H. *see* Roden, M.
- Tan, Y. *see* Ng, T.-P.
- Tang, Y.H. *see* Choi, S.W.
- Tarassenko, L. *see* Farmer, A.
- Targher, G., Bertolini, L., Scala, L., Zoppini, G., Zenari, L. & Falezza, G. Non-alcoholic hepatic steatosis and its relation to increased plasma biomarkers of inflammation and endothelial dysfunction in non-diabetic men. Role of visceral adipose tissue, 1354
- Targher, G., Bertolini, L., Zoppini, G., Zenari, L. & Falezza, G. Increased plasma markers of inflammation and endothelial dysfunction and their association with microvascular complications in Type 1 diabetic patients without clinically manifest macroangiopathy, 999
- Targher, G., Bertolini, L., Zoppini, G., Zenari, L. & Falezza, G. Relationship of non-alcoholic hepatic steatosis to cortisol secretion in diet-controlled Type 2 diabetic patients, 1146
- Tawata, M. *see* Kouki, T.
- Tay, E.-G. *see* Ng, T.-P.
- Tayebjee, M.H. *see* Tan, K.T.
- Tayebjee, M.H., Lip, G.Y.H. & MacFadyen, R.J. What role do extracellular matrix changes contribute to the cardiovascular disease burden of diabetes mellitus?, 1628
- Taylor, A. *see* Leese, G.P.
- Taylor, K.W. Viruses and diabetes, 957
- Taylor, R. *see* Advani, A.
- Taylor, R. *see* Al-Bermani, A.
- Teesalu, R. *see* Ainla, T.
- Temple, R.C. *see* Roland, J.M.
- Temple, R.C. *see* Sampson, M.J.
- Teramo, K. *see* Loukovaara, M.
- Thai, A.-C. *see* Ng, T.-P.
- Thayssen, P. *see* Hoffmann, S.K.S.
- Theodoropoulou, A. *see* Habeos, I.G.
- Thomas, B. *see* Min, Y.
- Thomas, M.C., Walker, M.K., Emberson, J.R., Thomson, A.G., Lawlor, D.A., Ebrahim, S. & Whincup, P.H. Prevalence of undiagnosed Type 2 diabetes and impaired fasting glucose in older British men and women, 789
- Thomson, M.J., Biddulph, J.P., Cull, C.A. & Holman, R.R. Reporting of diabetes on death certificates using data from the UK Prospective Diabetes Study, 1031
- Thompson, C.F. *see* Simmons, D.
- Thompson, L., Nester, C., Stuart, L. & Wiles, P. Interclinician variation in diabetes foot assessment—a national lottery?, 196
- Thomson, A.G. *see* Thomas, M.C.
- Thorsby, P.M. *see* Aas, A.M.
- Thorsson, A.V. *see* Tryggvason, G.
- Thorsteinsson, B. *see* Høi-Hansen, T.
- Tian, H. *see* Ran, X.
- Tian, H. *see* Wang, X.L.
- Timms, P.M. *see* Boucher, B.J.
- Timpson, N. *see* Lawlor, D.A.
- Tinahones, F. *see* Morcillo, S.
- Tochikubo, O. *see* Yasuda, G.
- Todd, I. *see* Damanhour, L.H.
- Tognoni, G. *see* Franciosi, M.
- van Tol, A. *see* Oomen, P.H.N.
- Tolias, A. *see* Meyer, C.
- Tomasi, J.P. *see* Buysschaert, M.
- Tong, N. *see* Ran, X.
- Tonozuka, N. *see* Matsumura, Y.
- Tooke, J.E. *see* Fegan, P.G.
- Tooke, J.E. *see* Gooding, K.M.
- Torffvit, O., Lövestam-Adrian, M., Agardh, E. & Agardh, C.-D. Nephropathy, but not retinopathy, is associated with the development of heart disease in Type 1 diabetes: a 12-year observation study of 462 patients, 723
- Torppa, J. *see* Pajunen, P.
- Toth, E.L. *see* Johnson, J.A.
- Toubro, S. *see* Black, E.
- Treece, K. *see* Pound, N.
- Treece, K.A. *see* Chipchase, S.Y.
- Tregnaghi, C. *see* Giannarelli, R.
- Tringham, J.R. *see* Skinner, T.C.
- Tringham, J.R., Khunti, K.K., Davies, M.J. Reply, 1286
- Tromp-Wever, A.M.E. *see* van der Ven, N.C.W.
- Tryggvason, G., Indridason, O.S., Thorsson, A.V., Hreidarsson, A.B. & Palsson, R. Unchanged incidence of diabetic nephropathy in Type 1 diabetes: a nation-wide study in Iceland, 182

- Tsai, S.-T. *see* Chen, L.-K.  
 Tschiedel, B. *see* Crispim, D.  
 Tseng, C.-H., Chong, C.-K., Sheu, J.-J., Wu, T.-H. & Tseng, C.-P. Prevalence and risk factors for stroke in Type 2 diabetic patients in Taiwan: a cross-sectional survey of a national sample by telephone interview, 477  
 Tseng, C.-P. *see* Tseng, C.-H.  
 Tsugane, S. for the JPHC Study Group *see* Waki, K.  
 Tsuji, A. *see* Matsumura, Y.  
 Tsuzura, S. *see* Ikeda, Y.  
 Tubiana-Rufi, N. *see* Du Pasquier-Fediaevsky, L.  
 Tuomilehto, J. *see* Bosnyak, Z.  
 Tuomilehto, J. *see* Dong, Y.  
 Tuomilehto, J. *see* Pajunen, P.  
 Tuomilehto, J. *see* Söderberg, S.  
 Tuomilehto, J. *see* Ylihäsälä, H.  
 Turco, S. *see* Di Bonito, P.  
 Tuxen, C. *see* Hildebrandt, P.  
 Twetman, S., Petersson, G.H. & Bratthall, D. Caries risk assessment as a predictor of metabolic control in young Type 1 diabetics, 312  
 Twisk, J.W.R. *see* van der Ven, N.C.W.  
  
 Uccioli, L. *see* Caselli, A.  
 Uccioli, L. *see* D'Ambrogio, E.  
 Uccioli, L. *see* Faglia, E.  
 Ukkola, O. *see* Kunnari, A.  
 Umemura, S. *see* Yasuda, G.  
 Unwin, N.C. *see* Adams, J.  
 Urhammer, S.A. *see* Bing, C.  
 Urhammer, S.A. *see* Hamid, Y.H.  
 Urquhart, R. *see* Roden, M.  
  
 Vaag, A. *see* Koster, B.  
 Vaaler, S. *see* Hussain, A.  
 Vagenakis, A.G. *see* Habeos, I.G.  
 Valabhji, J. *see* George, P.M.  
 Valensi, P. *see* N'Guyen Thi, N.  
 Valentine, W.J. *see* Roze, S.  
 Valentini, M. *see* Franciosi, M.  
 Valle, T.T. *see* Ylihäsälä, H.  
 Vámosi, I. *see* Barkai, L.  
 Varillas, V.F. *see* Boronat, M.  
 Várkonyi, T.T. *see* Nemes, A.  
 Varughese, G.I. *see* Buch, H.N.  
 Vazic Babic, V. *see* Mlinaric Missoni, E.  
 Veldhuis, J.D. *see* Hollingdal, M.  
 Veldman, B.A.J., Vervoort, G., Blom, H. & Smits, P. Reduced plasma total homocysteine concentrations in Type 1 diabetes mellitus is determined by increased renal clearance, 301  
 Velmurugan, K. *see* Mohan, V.  
 van der Ven, N.C.W., Hogenelst, M.H.E., Tromp-Wever, A.M.E., Twisk, J.W.R., van der Ploeg, H.M., Heine, R.J. & Snoek, F.J. Short-term effects of cognitive behavioural group training (CBGT) in adult Type 1 diabetes patients in prolonged poor glycaemic control. A randomized controlled trial, 1619  
 Vendrell, J., Broch, M., Fernandez-Real, J.M., Gutiérrez, C., Simón, I., Megia, A., Gallart, L., Ricart, W. & Richart, C. Tumour necrosis factor receptors (TNFRs) in Type 2 diabetes. Analysis of soluble plasma fractions and genetic variations of TNFR2 gene in a case-control study, 387  
 Verge, C.F. *see* Donaghue, K.C.  
 Vèrier-Mine, O., Chaturvedi, N., Webb, D. & Fuller, J.H. and The EURODIAB Prospective Complications Study Group. Is pregnancy a risk factor for microvascular complications? The EURODIAB Prospective Complications Study, 1503  
 Vervoort, G. *see* Veldman, B.A.J.  
 Vida, J.M. *see* Paniagua, J.A.  
 Vijan, S., Stuart, N.S., Fitzgerald, J.T., Ronis, D.L., Hayward, R.A., Slater, S. & Hofer, T.P. Barriers to following dietary recommendations in Type 2 diabetes, 32  
 Villegas, R. *see* Perry, I.J.  
 Vimalaswaran, K.S., Radha, V., Ghosh, S., Majumder, P.P., Deepa, R., Babu, H.N.S., Rao, M.R.S. & Mohan, V. Peroxisome proliferator-activated receptor- $\gamma$  co-activator-1 $\alpha$  (PGC-1 $\alpha$ ) gene polymorphisms and their relationship to Type 2 diabetes in Asian Indians, 1516  
 Vissing, H. *see* Hamid, Y.H.  
 Vitali, L. *see* Perriello, G.  
 Vlachos, L. *see* Meyer, C.  
 Volpato, S. *see* Blaum, C.S.  
 Volpe, L. *see* Di Cianni, G.  
 Vuagnat, H. *see* Pataky, Z.  
 Vukelic, M. *see* Mlinaric Missoni, E.  
 Vuorenmaa, T. *see* Pajunen, P.  
  
 Wachtell, K. *see* Hildebrandt, P.  
 Wagner, Z. *see* Szelestey, T.  
 Wainstein, J., Metzger, M., Boaz, M., Minuchin, O., Cohen, Y., Yaffe, A., Yerushalmy, Y., Raz, I. & Harman-Boehm, I. Insulin pump therapy vs. multiple daily injections in obese Type 2 diabetic patients, 1037  
 Wakamatsu, T. *see* Jimba, S.  
 Waki, K., Noda, M., Sasaki, S., Matsumura, Y., Takahashi, Y., Isogawa, A., Ohashi, Y., Kadowaki, T. & Tsugane, S. for the JPHC Study Group Alcohol consumption and other risk factors for self-reported diabetes among middle-aged Japanese: a population-based prospective study in the JPHC study cohort I, 323  
 Walker, A.B. *see* Buch, H.N.  
 Walker, J. The kidney and hypertension in diabetes mellitus, 6th Edition, 960  
 Walker, M. *see* Bennett, S.M.A.  
 Walker, M. *see* Zeggini, E.  
 Walker, M.K. *see* Thomas, M.C.  
 Wallentin, L. *see* Dotevall, A.  
 Wallymahmed, M. *see* Woodward, A.  
 Wallymahmed, M.E., Dawes, S., Clarke, G., Saunders, S., Younis, N. & MacFarlane, I.A. Hospital in-patients with diabetes: increasing prevalence and management problems, 107  
 Wang, C. *see* Ran, X.  
 Wang, H. *see* Ran, X.  
 Wang, X.L., Lu, J.M., Pan, C.Y., Tian, H. & Li, C.L. A comparison of urinary albumin excretion rate and microalbuminuria in various glucose tolerance subjects, 332  
 Wang, Y. *see* Dong, Y.  
 Ward, P. *see* Betts, P.  
 Waris, M.E., Koskinen, J.O., Simell, O., Knip, M., Ilonen, J. Onset of  $\beta$ -cell autoimmunity is not associated with elevated concentration of C-reactive protein in children at genetic risk for Type 1 diabetes, 1123  
 Warren, R.E. *see* Graveling, A.J.



- the Warren 3/UK GoKind Study Group, 69
- Wasada, T. *see* Jimba, S.
- Wäscher, I. *see* Sanchez-Albisua, I.
- Watanabe, H. *see* Yamanouchi, T.
- Watts, G.F. *see* Dogra, G.K.
- Watts, G.F. *see* Keech, A.C.
- Waugh, N.R. *see* Royle, P.L.
- Webb, D. *see* Vèrier-Mine, O.
- Weisbrod, C.J., Eastwood, P.R., O'Driscoll, G. & Green, D.J.  
Abnormal ventilatory responses to hypoxia in Type 2 diabetes, 563
- Wellard, R.M. *see* Cameron, F.J.
- Welsman, J.R. *see* Middlebrooke, A.R.
- Wen, J. *see* Guang-Da, X.
- Wentzel-Larsen, T. *see* Graue, M.
- Werther, G.A. *see* Cameron, F.J.
- Werther, G.A. *see* Northam, E.A.
- Whincup, P.H. *see* Thomas, M.C.
- Whincup, P.H., Gilg, J.A., Owen, C.G., Odoki, K., Alberti, K.G.M.M. & Cook, D.G. British South Asians aged 13–16 years have higher fasting glucose and insulin levels than Europeans, 1275
- Whitcomb, D.C. *see* Schneider, A.
- White, M. *see* Adams, J.
- Widel, M. *see* Roden, M.
- Wild, S.H., Smith, F.B., Lee, A.J. & Fowkes, F.G.R. Criteria for previously undiagnosed diabetes and risk of mortality: 15-year follow-up of the Edinburgh Artery Study cohort, 490
- Wilding, J. *see* Woodward, A.
- Wilding, J.P.H. *see* Bennett, S.M.A.
- Wiles, P. *see* Thompson, L.
- Wilkin, T. *see* Lüders, S.
- Wilkin, T. *see* Betts, P.
- Wilkin, T. Is the 'Accelerator Hypothesis' worthy of our attention? 1458
- Wilson, C., Curtis, J., Lipke, S., Bochenski, C. & Gilliland, S. Nurse case manager effectiveness and case load in a large clinical practice: implications for workforce development, 1116
- Wittmann, I. *see* Szelestei, T.
- Wittrup, M. *see* Bodtger, U.
- Wittrup-Jensen, K.U. *see* Lee, A.J.
- Wolf, J. *see* Sanchez-Albisua, I.
- Woo, G.C. *see* Choi, S.W.
- Woodward, A., Wallymahmed, M., Wilding, J. & Gill, G. Improved glycaemic control—an unintended benefit of a nurse-led cardiovascular risk reduction clinic, 1272
- Worman, H.J. *see* Lüdtkke, A.
- Wotherspoon, F., Browne, D.L., Meeking, D.R., Allard, S.E., Munday, L.J., Shaw, K.M. & Cummings, M.H. The contribution of nitric oxide and vasodilatory prostanoids to bradykinin-mediated vasodilation in Type 1 diabetes, 697
- Wouterse, A.C. *see* Bisseling, T.M.
- Wraight, P.R., Lawrence, S.M., Campbell, D.A. & Colman, P.G. Creation of a multidisciplinary, evidence based, clinical guideline for the assessment, investigation and management of acute diabetes related foot complications, 127
- Wu, L.-Y. *see* Hung, Y.-J.
- Wu, T.-H. *see* Tseng, C.-H.
- Wulffélé, M.G., Kooy, A., Leher, P., Bets, D., Donker, A.J.M. & Stehouwer, C.D.A. Does metformin decrease blood pressure in patients with Type 2 diabetes intensively treated with insulin?, 907
- Wyatt, J.W., Frias, J.L., Hoyme, H.E., Jovanovic, L., Kaaia, R., Brown, F., Garg, S., Lee-Parritz, A., Seely, E.W., Kerr, L., Mattoo, V., Tan, M. & the IONS study group. Congenital anomaly rate in offspring of mothers with diabetes treated with insulin lispro during pregnancy, 803
- Xue, Q.L. *see* Blaum, C.S.
- Yaffe, A. *see* Wainstein, J.
- Yamada, M. *see* Shimizu, H.
- Yamamoto, S. *see* Yoneda, Y.
- Yamamoto, Y. *see* Nishi, M.
- Yamanouchi, T., Sakai, T., Igarashi, K., Ichianagi, K., Watanabe, H. & Kawasaki, T. Comparison of metabolic effects of pioglitazone, metformin, and glimepiride over 1 year in Japanese patients with newly diagnosed Type 2 diabetes, 980
- Yan, W. *see* Chen, S.
- Yao, Z. *see* Chen, S.
- Yasuda, G., Hasegawa, K., Kuji, T., Ogawa, N., Shimura, G., Umemura, S. & Tochikubo, O. Effects of doxazosin on ambulatory blood pressure and sympathetic nervous activity in hypertensive Type 2 diabetic patients with overt nephropathy, 1394
- Yasuda, H. *see* Sakuta, H.
- Yen, C.-J. *see* Jiang, Y.-D.
- Yerushalmi, Y. *see* Wainstein, J.
- Yeung, G.S.P. *see* Choi, S.W.
- Yeung, V.T.F. *see* Choi, S.W.
- Yiu, C.H. *see* Choi, S.W.
- Yliharsila, H., Lindström, J., Eriksson, J.G., Jousilahti, P., Valle, T.T., Sundvall, J. & Tuomilehto, J. Prevalence of diabetes and impaired glucose regulation in 45- to 64-year-old individuals in three areas of Finland, 88
- Yoneda, Y. & Yamamoto, S. Cerebral cortical laminar necrosis on diffusion-weighted MRI in hypoglycaemic encephalopathy, 1098
- Yoo, S.-J. *see* Ko, S.-H.
- Yoon, K.H. *see* Moon, M.K.
- Yoshida, H. *see* Nagaya, T.
- Yoshikawa, T. *see* Fukui, M.
- Youn, B.S. *see* Lee, Y.Y.
- Young, I.S. *see* McSorley, P.T.
- Young, R.J. *see* McElduff, P.
- Younis, N. *see* Wallymahmed, M.E.
- Yu, H. *see* Dong, Y.
- Yudkin, J.S. *see* Oswald, G.
- Zahlmann, G. *see* Schneider, S.
- Zaki, M. *see* Abdella, N.A.
- Zakrzewska, K.E. *see* Roze, S.
- Zeggini, E., Parkinson, J.R.C., Halford, S., Owen, K.R., Walker, M., Hitman, G.A., Levy, J.C., Sampson, M.J., Frayling, T.M., Hattersley, A.T. & McCarthy, M.I. Examining the relationships between the Pro12Ala variant in *PPARG* and Type 2 diabetes-related traits in UK samples, 1696
- Zenari, L. *see* Targher, G.
- Zhang, X. *see* Armour, T.A.
- Zhao, T. *see* Ran, X.
- Zimmet, P. *see* Söderberg, S.
- Zinman, B. *see* Pollex, R.L.
- Zoppini, G. *see* Targher, G.

## Subject Index

- 24-h ambulatory blood pressure monitoring, 907
- ABCC8, 590
- $\alpha_1$ -adrenergic antagonists, 1394
- $\alpha$ -glucosidase inhibition, 470
- Aborigines, 1589
- acarbose, 470
- accelerator hypothesis, 144
- accident and emergency department, 1766
- Achilles tendon, 1713
- activity monitor, 893
- acute complications, 1636
- acute coronary syndromes, 243, 1542
- acute medical admission, 937
- acute myocardial infarction, 1321
- adherence to diabetes regimen, 427
- adherence, 1379
- adipokines, 674
- adiponectin, 569, 1024
- adolescent, 1522
- adolescents, 92, 427, 893
- ADP-ribosyl cyclase, 1770
- adult, 88
- advanced glycation endproducts, 1218
- aerobic fitness, 483
- affective disorder, 293
- Africa, 1642
- African-Brazilians, 1683
- age, 1476
- AIDS, 218
- albuminuria, 723
- Alcohol consumption, 1446
- alcohol, 243, 1359
- aldose reductase, 1315
- allergy, 102
- ambulatory blood pressure measurement, 1054
- ambulatory blood pressure monitoring, 1218, 1394
- American Indian/Alaska Native, 1116
- amputation, 1258, 1306
- angioplasty, 460
- angiotensin-converting enzyme gene polymorphism, 1054
- angiotensin-converting enzyme, 1167
- aniridia, 641
- ankle-brachial index, 1310
- anomalies, 803
- anti-diabetics, 2
- antioxidant, 703
- antioxidants, 1347
- anxiety, 1497
- anxiety disorder, 293
- ApoE polymorphism, 21
- apolipoprotein AV, 1690
- aripiprazole, 1440
- ascorbic acid, 1347
- Asian Indians, 863, 1151, 1516
- aspart, 606
- association study, 946
- association, 1696
- Atherosclerosis, 1212
- atherosclerosis, 81, 1628, 1657, 1701
- atypical antipsychotics, 1440
- autoantibodies, 340, 448, 1770
- auto-immunity, 889
- autoimmunity, 1770
- autonomic neuropathy, 1614
- $\beta$ -cell function, 1744
- $\beta$ -cell, 1016
- $\beta$ -hydroxybutyrate, 221
- Bangladesh, 931
- barriers, 32
- basal insulin delivery, 382
- behaviour models, 415
- behavioural intervention, 1619
- beliefs, 1497
- bereavement, 1338
- beta-cell function, 1677
- biomechanics, 1713
- biphasic insulin aspart, 273
- birth order, 200
- birth weight, 21, 200
- birthweight, 440, 921
- bite injury, 631
- blood glucose self-monitoring, 1372
- blood glucose, 1326
- blood pressure, 882, 907, 1272, 1589
- blood-brain barrier, 1278
- BMI, 921
- BNP, 188
- body fat, 863
- body mass index, 931, 1107, 1482, 1696
- bodyweight, 850
- body weight, 1574
- body weight changes, 316
- bone mineral density, 756
- bradykinin, 697
- brain, 648
- breast feeding, 177
- breathing pattern, 563
- bupropion, 658
- Caesarean section, 177
- candidaemia, 1252
- capillary blood glucose, 1173
- capillary glucose, 599
- capillary pressure, 536
- carbohydrate-rich meals, 213
- cardiac neuropathy, 1072
- cardiomyopathy, 1218
- cardiovascular defects and gestational diabetes, 775
- cardiovascular disease, 26, 256, 306, 950, 1031, 1326, 1558
- cardiovascular diseases, 118
- cardiovascular mortality, 723
- cardiovascular risk factors, 551, 612, 1737
- caries, 312
- carotid artery, 81

- carotid atherosclerosis, 1446  
 carpal tunnel syndrome, 625  
 Carter effect, 1551  
 case management, 1116  
 case-control study, 775  
 Caucasian-Brazilians, 1683  
 cause-specific mortality, 1636  
 cerebral oedema, 645  
 cerebrovascular disease, 583  
 Charcot foot, 1707  
 Charcot neuropathic osteoarthropathy, 756  
 chemoreflex, 563  
 childhood, 648, 921  
 children and adolescents, 1054  
 children, 92, 144, 312, 427, 711, 1566  
 Chinese, 1737  
 chronic complications, 1047  
 chronic critical limb ischaemia, 460  
 chronic diseases, 415  
 chronic painful diabetic neuropathy, 393  
 chronic pancreatitis, 744  
 cigarette smoking, 658  
 cirrhosis, 1129, 1530  
 claudication, 1310  
 clinical foot examination, 1310  
 clinical guidelines, 554  
 clinician variation, 196  
 coeliac disease, 889, 1079  
 cohort studies, 61, 1178  
 cohort study, 950  
 cohort, 1338  
 co-morbidity, 258  
 comorbidity, 293  
 complement C3, 1282  
 complications, 1451  
 congenital abnormalities, 1774  
 congestive heart failure, 986  
 continuous glucose monitoring system, 858  
 continuous subcutaneous insulin infusion, 382, 842, 1239  
 contraindications, 842  
 coronary artery disease, 258, 863, 1206, 1628  
 coronary flow reserve, 1614  
 coronary heart disease, 164  
 coronary microcirculation, 158  
 coronary vasoreactivity, 45  
 cortisol secretion, 1146  
 cost effective, 1239  
 costs, 1239  
 C-peptide, 1178, 1414  
 C-reactive protein, 278, 863, 1282  
 criteria, 1731  
 cross-cultural comparisons, 942  
 cross-national, 1379  
 cross-sectional studies, 336  
 cumulative incidence, 182  
 CURES, 863  
 cyclo-oxygenase, 503  
 cytomegalovirus, 218  
 death certificate, 1031  
 deconvolution analysis, 1408  
 dehydroepiandrosterone, 1446  
 delivery of care, 619  
 depression, 942, 1465  
 deprivation, 1364  
 Diabetes, 1005  
 diabetes, 32, 56, 107, 127, 306, 312, 332, 344, 448, 483, 503, 543, 563, 730, 749, 803, 840, 889, 950, 994, 1072, 1083, 1112, 1129, 1134, 1234, 1321, 1326, 1338, 1359, 1372, 1420, 1465, 1482, 1581, 1611, 1657, 1707, 1713, 1726  
 diabetes complications, 1060, 1258, 1306  
 diabetes knowledge, 427  
 diabetes mellitus, 61, 81, 88, 118, 144, 213, 218, 243, 249, 286, 323, 340, 348, 410, 497, 576, 619, 658, 768, 833, 871, 986, 1107, 1116, 1151, 1167, 1199, 1226, 1252, 1295, 1334, 1530, 1542, 1720, 1757  
 diabetes mellitus Type 2, 674  
 diabetes reporting, 1031  
 diabetic cardiomyopathy, 1720  
 diabetic complications, 348, 410  
 diabetic dyslipidaemia, 26  
 diabetic foot, 127, 460, 631, 762, 1487  
 diabetic ketoacidosis, 218, 466, 645, 842, 1440  
 diabetic nephropathy, 14, 182, 453, 1167, 1394, 1761, 762  
 diabetic neuropathy, 1608, 1763  
 diabetic polyneuropathy, 231  
 diabetic retinopathy, 794, 840  
 diagnosis, 102, 336, 833, 1173, 1434, 1731, 1757  
 diagnostic criteria, 490  
 diagnostic screening, 1012  
 diastolic function, 730  
 diet therapy, 316  
 diet, 821  
 dietary adherence, 32  
 dipeptidyl peptidase IV, 654  
 DKA, 221  
 DNA damage, 1347  
 DNA polymorphism, 1167  
 dose-response, 1016  
 dose-response relationship, 1107  
 dynamometer, 1401  
 early-onset diabetes mellitus, 641  
 echocardiography, 188, 1218, 1614  
 education, 1060  
 electrical spinal cord stimulation, 393  
 emergency department, 221  
 encephalopathy, 1098  
 endothelial dysfunction, 249, 999, 1212, 1354, 1663  
 endothelial function, 239, 697  
 endothelium, 503, 703, 1487, 1647, 1670  
 end-stage renal disease, 182  
 energy intake, 893  
 entrapment neuropathy, 625  
 epidemiology, 808, 336, 477, 775, 1199, 1263, 1420, 1551, 1751  
 erythrocyte, 914  
 essential fatty acids, 914  
 essential hypertension, 1394  
 ethnicity, 1226, 1343, 1598  
 etogenic, 821  
 evaluation, 1234  
 evidence-based information, 688  
 evidence-based medicine, 127  
 exercise, 316, 1581  
 experimental diabetes, 359  
 extra cellular matrix, 1628  
 familial partial lipodystrophy, 1611  
 family history, 207  
 family interventions, 1295  
 family practice, 336  
 fasting plasma glucose, 1234, 1275  
 fatigue, 1401  
 ferritin, 97  
 fetal insulin, 950  
 fetus, 434, 914  
 first-degree relative, 1551  
 first-phase insulin secretion, 1408  
 flow cytometry, 1657  
 foetoplacental circulation, 503  
 folic acid, 1774  
 follow-up study, 1107  
 foot health assessment, 196  
 foot pressure, 1487  
 foot sole, 762  
 foot ulcer, 1060, 1258, 1306  
 Framingham equation, 554  
 free fatty acids, 980  
 $\gamma$ -glutamyl transferase, 1359  
 $\gamma$ -glutamyltransferase, 1134  
 G protein-coupled receptor 40, 74  
 gastric bypass, 1761  
 general practice, 1173, 1246  
 genetic epidemiology, 946  
 genetic predisposition, 1167, 69  
 genetics, 74, 517, 590, 1516  
 gestational diabetes mellitus, 914, 1012, 1434, 1574  
 gestational diabetes, 177, 434, 775, 1267, 1536, 1731  
 gestational glucose intolerance, 1536  
 Ghrelin, 1157  
 GIP, 470  
 glargine, 374  
 gliclazide, 399, 973

- glimepiride, 980  
 GLP-1 (7-36 amide), 470  
 GLP-1, 654, 1016  
 glucokinase, 994  
 glucokinase regulatory protein, 1677  
 gluconeogenesis and insulin signalling, 2  
 glucose, 177, 207  
 glucose control, 882  
 glucose intolerance, 1091, 1206, 1212  
 glucose metabolism, 453  
 glucose production, 453  
 glucose tolerance, 21  
 glucose transport, 946, 1278  
 glutamic acid decarboxylase, 448  
 gluten-free diet, 1079  
 glycaemia, 980  
 glycaemic control, 344, 963, 1060, 1343, 1347, 1364, 1372, 1598  
 glycated haemoglobin, 207, 833  
 glycogen metabolism, 2  
 GPR40, 74  
 Greece, 1581  
 growth, 1079  
 guidelines, 32, 127, 937  
  
 habit, 415  
 haemoglobin A<sub>1c</sub>, 612  
 HbA<sub>1c</sub>, 344, 427, 1079, 1272, 1566, 1726  
 health care, 711  
 health outcomes, 1116  
 health services use, 1047  
 health services, 1246  
 healthcare professionals, 1246  
 health-related quality of life, 1522  
 heart, 730  
 heart disease, 723  
 heart-rate variability, 1072  
 heel, 1258  
 height, 921  
 hepatic steatosis, 1146, 1354  
 hepatitis C, 1530  
 hepatitis C virus, 340  
 high-protein, 821  
 HIV infection, 218  
 HLA, 137  
 HNF-1 $\alpha$ , 137  
 HNF-4 $\alpha$  genes, 137  
 HOMA insulin resistance, 1476  
 HOMA-%S index, 39  
 HOMA-IR, 346  
 homocysteine, 1359, 301  
 hospital in-patients, 107  
 hyperglycaemia, 152, 45, 470, 543, 768, 871, 1185, 1252, 1321, 1401, 1440  
 hyperglycaemic hyperosmolar syndrome in children, 645  
 hyperinsulinaemia, 118, 768  
 hyperinsulinaemic glucose clamps, 1064  
 hyperlipidaemia, 1611  
 hypertension, 188, 737, 1005, 1267, 1605  
  
 hypertriglyceridaemia, 1690  
 hypertrophy, 730, 1005  
 hypoaldosteronism, 1605  
 hypoglycaemia, 92, 152, 749, 842, 850, 858, 1444  
 Hypoglycaemia, 1098  
 hypoglycaemic agents, 986  
  
 IA-2, 448  
 IFG, 1476  
 IGT, 1476  
 IL-10, 719  
 IL-8, 719  
 illness perceptions, 286  
 impact, 1497  
 impaired fasting glucose, 61, 599, 789, 1141, 1766  
 impaired fasting glycaemia, 332  
 impaired glucose tolerance, 61, 332, 599  
 incidence, 348  
 incidental hyperglycaemia, 937  
 incretin hormones, 470  
 in-depth interview, 286  
 indications, 842  
 Indigenous populations, 833  
 indomethacin, 503  
 inflammation, 1212, 1701  
 information systems, 1226  
 insulin, 316, 634, 749, 1037, 1060, 1178, 1239, 1414, 1434, 1598, 1677  
 insulin action, 1064  
 insulin aspart, 850  
 insulin detemir, 850  
 insulin discontinuation, 1744  
 insulin glargine, 382, 1444  
 insulin lispro, 803  
 insulin pump therapy, 92  
 insulin resistance, 39, 56, 266, 306, 346, 517, 569, 674, 703, 808, 921, 1091, 1157, 1190, 1206, 1275, 1589, 1611, 1614, 1701, 1720, 1751  
 insulin resistance syndrome, 39  
 insulin secretion, 52, 406, 470, 946  
 Insulin sensitivity, 1101  
 insulin sensitivity, 1024, 52, 1574  
 insulin therapy, 118, 177, 877  
 Insulin, 102  
 insulin-dependent, 118  
 insulin-like growth factor binding protein-1, 434  
 insulin-treated diabetes, 440  
 intensive management, 410  
 Inter99, 74  
 intermittent claudication, 1608  
 Internet, 688  
 intervention, 1522  
 intima media thickness, 1558  
 intravenous insulin, 221  
 iontophoresis, 536, 1670  
 ischaemia, 231  
  
 islet antibodies, 137  
 isotopes, 453  
 Italy, 1263  
  
 K<sub>ITT</sub> index, 39  
 KCNJ11, 590  
 ketoacidosis, 1414  
 ketosis-prone diabetes, 1744  
 kidney, 453  
 Korea, 1134  
 Korean population, 569  
 Korean, 1161  
 lactic acidosis, 1451  
 laminar necrosis, 1098  
 laser photocoagulation therapy, 1083  
 left ventricular function, 1005  
 left ventricular hypertrophy, 188, 1218  
 leptin, 278  
 life table analysis, 1551  
 lifetime risk, 625  
 lipids, 1670  
 lispro mixture, 374  
 literature searching, 1386  
 liver enzymes, 973  
 longitudinal cohort study, 427  
 long-term, 399  
 long-term effectiveness, 393  
 long-term storage, 1726  
 low-carbohydrate, 821  
 low-carbohydrate diet, 1605  
  
 macrosomia, 440, 1536  
 macrovascular, 348  
 macrovascular disease, 164, 882  
 magnetic resonance, 648  
 malformations, 803  
 malformations and BMI, 775  
 malnutrition, 1072  
 management, 107  
 mass screening, 794  
 maternal age, 200  
 maternal diabetes, 693  
 maternally inherited diabetes and deafness, 1683  
 matrix metalloproteinases, 1628  
 maturity-onset diabetes of the young, 1012  
 Mauritius, 61  
 maximal muscle strength, 1401  
 maximum hyperaemia, 536  
 MCP-1, 719  
 meal-time glycaemic control, 606  
 mechanisms of  $\beta$ -cell dysfunction, 963  
 Mediterranean diet, 1581  
 men, 1446  
 meta-analysis, 590, 1295, 1386  
 metabolic control, 427, 900, 1522  
 metabolic syndrome, 266, 808, 1134, 1146, 1157, 1206, 1354, 1465, 1510, 1589, 1751  
 metabolism, 648

- metformin, 346, 497, 634, 654, 907, 973, 980, 1101, 1282, 1451
- microalbuminuria, 14, 239, 332, 711, 1647
- microdialysis, 56
- microvascular, 348, 483, 1670
- microvascular complications, 256, 999, 1503
- microvascular disease, 882, 1487
- microvascular filtration capacity, 536
- middle age, 1199
- milk consumption, 808
- minute ventilation, 563
- mitochondrial G1888A variant, 1683
- modelling, 1239
- MODY, 517
- MODY3, 406
- monitoring, 937
- morbidity and mortality, 497
- mortality, 258, 466, 490, 543, 1031, 1252, 1321, 1642
- mouse, 359
- MRI, 1098
- MspI polymorphism of the APO AI gene, 782
- multi-factorial intervention, 249
- multiple daily injection, 1239
- muscle infarction, 1757
- mutation, 994
- myocardial infarction, 1334, 1542
- myocardial perfusion, 45
- myotonic dystrophy, 346
- nateglinide, 158
- neonatal deaths, 440
- neonatal Ponderal Index, 1536
- nephropathy, 69, 387
- NEUROD1, 1012
- neuropathy, 756, 1258, 1306, 1315, 1707
- newly diagnosed type 2 diabetes, 1310
- newly diagnosed Type 2 diabetes, 1737
- nicotine replacement therapy, 658
- nitric oxide, 697
- nitric oxide synthase, 1663
- nitrotyrosine, 1185
- nocturnal pain, 1763
- non-alcoholic fatty liver disease, 1129, 1141
- non-insulin treated, 900
- non-insulin-dependent, 118
- non-insulin-dependent diabetes mellitus, 316, 336, 1178
- non-prescribable medicine, 1492
- non-synonymous substitution, 1690
- North American Aboriginal population, 1510
- nurse-led clinics, 1272
- nursing, 1116
- obese, 1037
- obesity, 52, 144, 278, 612, 1107, 1129, 1161, 1199, 1476, 1482, 1737, 1761
- olanzapine, 1440
- omega-3 polyunsaturated fatty acids, 1465
- ophthalmology, 1112
- optimization, 382
- oral agents, 374
- oral glucose tolerance test, 88
- oral hypoglycaemic agents, 1774
- orlistat, 344, 612, 1737
- osteoarthropathy, 1707
- outcome, 1566
- outcome measures, 1306
- outcome study, 900
- oxidant-antioxidant indices, 1343
- oxidative stress, 1151, 1647
- oximetry, 460
- pain, 231
- pancreatic secretory trypsin inhibitor, 744
- parity, 1574
- paternal age, 200
- pathogenesis, 231
- patient education, 410
- patient preferences, 32
- patient satisfaction, 1246
- patient-centred care, 683
- PAX6, 641
- peripheral arterial disease, 1258, 1310
- peripheral arterial occlusive disease, 1608
- peripheral neuropathy, 1763
- peripheral pulses, 1310
- peripheral vascular disease, 460
- peroxisome proliferator activated receptor- $\gamma$  co-activator-1, 1516
- peroxisome proliferator activated receptor- $\gamma$ , 1516
- peroxisome proliferator-activated receptor  $\gamma$ , 1161
- peroxisome proliferator-activated receptor  $\gamma$ 2, 1696
- pharmacoepidemiology, 986
- pharmacokinetics, 273, 1444
- philosophy of mind, 415
- phosphoglycerides, 914
- phosphoinositide 3-kinase, 737
- phosphoisoforn, 434
- phospholipids, 1465
- physical activity, 893, 1206, 1581
- PI 3-kinase, 674
- pioglitazone, 399, 973, 980, 1101
- plantar fascia, 1713
- plantar pressure, 762
- plasma cholesteryl ester transfer protein, 768
- plasma glucose, 1173
- plasma phospholipid transfer protein, 768
- platelet microparticles, 1657
- platelets, 1657
- pleiotropism, 239
- polymorphism, 569, 946, 1157, 1161, 1663, 1677
- positron emission tomography, 45, 158
- post-transfusion, 340
- postural hypotension, 1072
- power spectral analysis, 1394
- practice nurses, 683
- practice organization, 1364
- precipitating factors, 466
- prediction, 1263
- pregnancy, 21, 803, 1503, 1774
- pregnancy outcome, 1536
- prevalence, 61, 88, 107, 293, 477, 789, 889, 1267, 1427, 1731, 1751
- primary care, 619, 683, 1364
- primary healthcare utilization, 1083
- probability, 1199
- prognosis, 1326, 1530
- proinsulin, 1178
- proliferative diabetic retinopathy, 719
- propensity score, 497
- prospective, 306
- prospective study, 323
- prostaglandins, 697
- protamine, 102
- protective factors, 1091
- protein kinase B, 674
- proteinuria, 14, 182
- proteinuria resolution, 1761
- provider behaviour, 1379
- psychiatric morbidity, 152
- psychological well-being, 1379
- psychosocial aspects, 1619
- pulsatile insulin secretion, 1064, 1408
- pulse-wave analysis, 1701
- pump, 1037
- pupillometry, 1315
- quality assurance, 794
- quality indicators, 1566
- quality of care, 1226, 619
- quality of life, 92, 1482
- QUICKI, 1101
- race, 1636
- randomized controlled trial, 907
- rat, 359
- recurrent hypoglycaemia, 1278
- registry, 1420
- regulatory subunit, 737
- relatives, 703
- reliability, 1726
- renal clearance, 301
- renal damage, 258
- renal failure, 730, 1642
- repaglinide, 1408
- resistin, 583
- retinopathy, 711, 723, 1112



- risk assessment, 312  
 risk factor, 323, 554, 1420  
 risk factors, 477, 599, 1047, 1263, 1434, 1510  
 risk factors for coronary heart disease, 278  
 risk score, 1234  
 rodent, 359, 631  
 safety, 393  
 satisfactory management, 937  
 school absenteeism, 421  
 screening, 207, 256, 599, 833, 840, 1112, 1234, 1434, 1497, 1766  
 secretagogue, 1016  
 self-care behaviours, 427  
 self-care strategy, 286  
 self-monitored blood glucose, 858  
 self-monitoring of blood glucose, 900  
 senescence, 1151  
 serine protease inhibitor Kazal type 1, 744  
 Sibutramine, 1024  
 single nucleotide polymorphism, 1690  
 skeletal muscle, 56  
 skin, 1487  
 skin test, 102  
 smoking, 81  
 socio-economic inequalities, 1047  
 soluble transferrin receptors, 97  
 South Asian Ethnicity Adolescence, 1275  
 specific congenital abnormalities, 693  
 spinal stenosis, 1763  
 stable angina pectoris, 1326  
 statin, 1647  
 statins, 239  
 stillbirth, 1267  
 stillbirths, 440  
 stored whole blood, 1726  
 stress, 1338  
 stroke, 477, 551, 1451  
 subclinical disorders, 293  
 subclinical inflammation, 999, 1354  
 Sudan, 213  
 sulfatide, 1190  
 sulfated lactosylceramide, 1190  
 sulf-lac-cer, 1190  
 sulphonylurea, 406  
 surgical debridement, 460  
 systematic review, 1295  
 systematic reviews, 1386  
 Taiwan, 286  
 Tanzania, 631  
 TC/HDL ratio, 26  
 telemedicine, 794, 1372  
 Telomere, 1151  
 temperature, 231  
 TGF $\beta$ -inducible gene h3( $\beta$ ig-h3), 14  
 therapeutic adherence, 415  
 therapy, 634  
 thiazolidinedione, 399, 1101, 1611  
 thiazolidinediones, 963, 986  
 thrombogenesis, 249  
 tissue Doppler imaging, 1720  
 tissue inhibitor of matrix metalloproteinase, 1628  
 TNF system, 387  
 TNF2 gene polymorphisms, 387  
 tobacco, 658  
 Torres Strait Islanders, 1589  
 total contact cast, 1707  
 transcription factor, 641  
 transforming growth factor- $\beta$ 1, 69  
 transgenics, 359  
 treatment, 1757  
 triglycerides, 21, 877, 914  
 Trinidad and Tobago, 619  
 truncal adiposity, 144  
 type 2 diabetes, 569  
 type 2-diabetes, 470  
 Type 1 diabetes, 45, 69, 137, 152, 164, 182, 200, 218, 273, 382, 421, 427, 434, 453, 551, 625, 697, 723, 756, 858, 893, 999, 1503, 1522, 1551, 1619, 1636, 1642, 1770  
 Type 1 diabetes mellitus, 239, 301, 1079, 1239, 1566  
 Type 2 diabetes, 52, 97, 158, 177, 188, 207, 256, 258, 374, 387, 517, 554, 583, 590, 606, 612, 634, 654, 658, 683, 688, 703, 756, 789, 863, 877, 882, 900, 907, 931, 942, 963, 973, 1037, 1047, 1064, 1141, 1146, 1161, 1173, 1190, 1246, 1263, 1364, 1414, 1446, 1451, 1492, 1497, 1510, 1516, 1558, 1605, 1647, 1663, 1670, 1677, 1683, 1690, 1766, 1770, 1774  
 Type 2 diabetes mellitus, 39, 74, 278, 399, 477, 490, 737, 744, 782, 1282, 1427  
 ulceration, 127  
 ultrasonography, 81  
 undiagnosed, 1766  
 undiagnosed diabetes, 599  
 United Kingdom, 1031  
 urban and rural population, 931  
 urinary albumin excretion rate, 332  
 urine albumin, 1558  
 utility, 1482  
 validation, 554  
 vanadium compounds, 2  
 vascular dysfunction, 871  
 vascular endothelial growth factor, 871, 1278  
 vascular risk, 164  
 ventricular arrhythmias and myocardial infarction, 576  
 very low calorie diet, 52  
 visceral fat, 1354  
 visceral obesity, 266  
 vision-threatening retinopathy, 1083  
 Vitamin E, 703  
 vitamin A, 1185  
 vitamin C, 1185, 1347  
 vitamin E, 1185  
 vitreous fluid, 719  
 von Willebrand factor, 871  
 waist-to-hip ratio, 931  
 walker, 1763  
 weight, 344, 1016  
 weight gain, 634  
 weight reduction, 266  
 West Indies, 619  
 WHO criteria, 789  
 windlass mechanism, 1713  
 within-person variability, 850  
 women, 1542

